

ANGELA GIBNEY

Moduli spaces of curves occupy a distinguished position in algebraic geometry. As moduli spaces, they give insight into the study of smooth curves and their degenerations, and as special varieties, they have played a principal role as a prototype for moduli of higher dimensional objects [KSB88, Ale02, HM06, HKT06, HKT09, CGK09]. We have recently seen revolutionary advances in the minimal model program [BCHM10], and moduli of curves provide one of the chief concrete, non-trivial settings where this nuanced theory has been exhibited, and explored [HH09, HH13, AFSvdW16, AFS16a, AFS16b].

Here I very briefly describe my own research about moduli spaces of curves, arranged along the following four general themes:

- (1) **Moduli of curves from vector bundles of conformal blocks**
- (2) **Vector bundles of conformal blocks from moduli of curves**
- (3) **Generalizations of moduli spaces of curves and their properties**
- (4) **Mori theory on the moduli space of curves**

I also briefly describe my newer work, with Prakash Belkale, where we show the section ring for the pair $(\text{Bun}_{\text{SL}(r)}(C), \mathcal{D})$ is finitely generated, for \mathcal{D} the determinant of cohomology line bundle.

MODULI OF CURVES, CONES OF POSITIVE CYCLES, AND CONFORMAL BLOCKS

The space $M_{g,n}$ is an irreducible, quasi-projective variety, whose points correspond to the set of equivalence classes of smooth curves of genus g , marked by n labeled points. Smooth curves degenerate to singular ones, and so $M_{g,n}$ is not compact. By adding curves with at worst simple nodal singularities and finitely many automorphisms, one obtains a projective variety $\overline{M}_{g,n}$ [DM69, KM76]. The stack $\overline{\mathcal{M}}_{g,n}$, parametrizing flat families of stable curves, reflects the geometry of $\overline{M}_{g,n}$, while being in certain ways easier to study. Since $\overline{M}_{0,n}$ is a fine moduli space, these points of view are equivalent in genus zero.

Given a projective variety X , the cone of nef divisors $\text{Nef}(X)$, is an invariant that consists of effective divisors that nonnegatively intersect all curves on X . The nef cone is dual to the closed cone of effective curves (aka the Mori cone). If $f : X \rightarrow Y$ is a morphism from X to another projective variety Y , and if A is an ample divisor on Y , then f^*A is an element of $\text{Nef}(X)$. While $\text{Nef}(X)$ is often difficult to describe, even its *shape* and *location* with respect to the cone of effective divisors gives valuable information about morphisms admitted by the variety X . For example, we know that the nef and effective cones of \overline{M}_g only touch at the origin [GKM02]. This implies that there are no morphisms (with connected fibers) from \overline{M}_g to any lower dimensional projective variety, other than a point.

The pseudo-effective cone $\overline{\text{Eff}}_k(X)$ is the closure of the cone generated by classes of k -dimensional subvarieties on X . If X is a smooth variety, one can define higher codimension analogues of cones of nef divisors by taking $\text{Nef}^k(X)$ to be dual to $\overline{\text{Eff}}_k(X)$. Many properties held by these cones when $k = 1$ fail more generally. To more accurately capture properties of positive cycles of higher codimension, Fulger and Lehmann have introduced three sub-cones of the nef cone $\text{Nef}^m(X)$. The smallest of these, the Pliant cone $\text{Pl}^m(X) \subset \text{Nef}^m(X)$, is the closure of the cone generated by monomials in Schur classes of globally generated vector bundles on X .

The stacks $\overline{\mathcal{M}}_{g,n}$ admit vector bundles of conformal blocks, constructed using affine Lie algebras [TK88, TUY89, Fak12]. In case g is zero, the vector bundles are globally generated: First Chern classes are base point free, giving elements of the nef cone, (products of) higher Chern classes give elements in Fulger and Lehmann's Pliant Cone. The interplay between intersection theory on $\overline{\mathcal{M}}_{g,n}$ and properties of these vector bundles, highlights nontrivial links between algebraic geometry and representation theory.

1. MODULI OF CURVES FROM VECTOR BUNDLES OF CONFORMAL BLOCKS

Extensions of families to the boundary. In [BG16, Theorem 1.2], we show there is a flat family over $\overline{\mathcal{M}}_g$ whose fibers over smooth curves C , are moduli spaces $SU_C(r)$, parameterizing semistable vector bundles of rank r with trivial determinant on C . All the fibers of the family are normal, projective varieties, constructed using conformal blocks.

Identification of faces of the nef cone. First Chern classes of vector bundles of conformal blocks, which are base point free on $\overline{\mathcal{M}}_{0,n}$, have been shown to give rise to many known contraction maps [GJMS13, AGS14], including the tautological maps. These can also be used to study the nef cone for higher genus curves [Fak12]. For instance, in [Gib12], I identify one of the faces of the nef cone of $\overline{\mathcal{M}}_g$, showing it corresponds to extensions of the Torelli map from $\overline{\mathcal{M}}_g$ to certain types of compactifications of A_g , the moduli space of Abelian varieties of dimension g .

New birational models. The study of alternative compactifications of moduli spaces of pointed curves has been important both in understanding families of curves, as well as giving an explicit description of the birational geometry (in the sense of Mori theory) of $\overline{\mathcal{M}}_{g,n}$. Conformal blocks have given new tools for providing modular interpretations of images of birational contractions of $\overline{\mathcal{M}}_{0,n}$. These new moduli spaces have been shown to include for example, cyclic covers [Fed11], and GIT quotients [Gia13, GJM13, GJMS13] generalizing Kapranov’s compactifications of $\mathcal{M}_{0,n}$ [Kap93a, Kap93b]. The latter giving a different (but overlapping) set of modular interpretations than described in the work of Smyth [Smy13], in that they parametrize “embedded curves”, as opposed to abstract ones (cf. [GJM13, p.245]).

Full-dimensional subcones of the positive cones, and finite generation in codimension one. Building on work of [Fak12], in recent work [GM16], we exhibit a full dimensional subcone of the Pliant cone $\text{Pl}^k(\overline{\mathcal{M}}_{0,n})$ constructed from (products of) Chern classes of vector bundles of conformal blocks. In [GG12], we show the infinite set of type A , level one conformal blocks divisors on $\overline{\mathcal{M}}_{0,n}$ spans a finitely generated, full-dimensional subcone of the cone of nef divisors. In [Kaz14], my former postdoc A. Kazanova has shown that the infinite set of S_n invariant, rank one, conformal blocks divisors for \mathfrak{sl}_n on $\overline{\mathcal{M}}_{0,n}$, which she identifies, spans a finitely generated, full-dimensional cone generated by level one divisors. In [Hob15], my student N. Hobson has shown that infinite set of rank one, conformal blocks divisors for \mathfrak{sl}_2 (and their generalizations) on $\overline{\mathcal{M}}_{0,n}$, which she identifies, spans a finitely generated, full-dimensional cone generated by level one divisors.

2. VECTOR BUNDLES OF CONFORMAL BLOCKS FROM THE MODULI SPACE OF CURVES

Geometric interpretations at points on the boundary. At smooth curves, conformal blocks correspond to generalized theta functions [BL94, Fal94, KNR94, Pau96, LS97], and in [BG16, Theorems 1.3 and 1.4], we show this is true for singular curves as well, in prescribed circumstances. As we show in [BGK16], using intersection theory on $\overline{\mathcal{M}}_g$, this does not hold in all situations.

Vanishing and identities. Characteristic classes of vector bundles of conformal blocks are subject to a number of types of identities: In [BGM15, Fak12], new rank and level identities are shown to hold for first Chern classes at the critical level, and divisors are trivial above the critical level. In [BG16], we show Chern characters in type A are quasi-polynomial. In [BGM16] we show under certain hypothesis, first Chern classes are subject to additive identities.

Questions of nonvanishing. Inspired by unexpected examples of trivial divisors, in [BGM16] we introduce the problem of finding necessary and sufficient conditions to determine when the first Chern class of a bundle is nontrivial. Necessary and sufficient conditions for nonvanishing are found for \mathfrak{sl}_2 .

3. GENERALIZATIONS OF THE MODULI SPACE OF CURVES, AND ITS PROPERTIES

Pointed trees of projective spaces. In [CGK09], Chen, Krashen and I construct a space $T_{d,n}$, whose points correspond to n -pointed stable rooted trees of d -dimensional projective spaces, which for $d = 1$, are $(n + 1)$ -pointed stable rational curves. Smooth and projective, with boundary a smooth normal crossings divisor, the space $T_{d,n}$ has an inductive construction analogous to but differing from Keel's construction of $\overline{M}_{0,n}$. We describe its Chow groups, Chow ring, Chow motive, and Poincaré polynomials, and use this information to answer a question of Fulton and MacPherson.

Tropical compactifications and a generalization of the F-Conjecture. While generally not a toric variety, it is possible to embed $\overline{M}_{0,n}$ into a noncomplete toric variety X_Δ , and in [GM10], we give equations for $\overline{M}_{0,n}$ in the Cox ring of X_Δ . In [GM12], we study varieties X that, like $\overline{M}_{0,n}$, can be embedded in a toric variety X_Δ , as a tropical compactification. We use X_Δ to define cones which are upper and lower bounds for $\text{Nef}(X)$. For $X = \overline{M}_{0,n}$, we show that our upper bound cone $U(\overline{M}_{0,n})$, is the cone of divisors predicted by the F-Conjecture to be equal to $\text{Nef}(\overline{M}_{0,n})$. In other words, this result generalizes the F-Conjecture on $\overline{M}_{0,n}$ to varieties that like $\overline{M}_{0,n}$, are tropical compactifications.

4. MORI THEORY ON THE MODULI SPACE OF CURVES

The F-Conjecture asserts that a divisor on $\overline{M}_{g,n}$ is nef if and only if it nonnegatively intersects one of a finite number of rational curves on $\overline{M}_{g,n}$ called F-Curves. In particular, the F-Conjecture implies that $\text{Nef}(\overline{M}_{g,n})$ is polyhedral, the convex hull of a finite number of extremal rays.

In [GKM02], we proved that the F-Conjecture on $\overline{M}_{0,g+n}$ implies the F-Conjecture on $\overline{M}_{g,n}$. The conjecture holds for low g and n [KM13, GKM02, FG03, Gib09]. In [Gib09], I reduce the F-conjecture on $\overline{M}_{g,n}$ to showing that certain divisors in $\overline{M}_{0,N}$ for $N \leq g + n$ are equivalent to the sum of the canonical divisor plus an effective divisor supported on the boundary. As an application, I give numerical criteria, which if satisfied by a divisor D on \overline{M}_g , show that D is ample. Using a computer program, The Nef Wizard, written by Krashen, this is used to verify the conjecture for $g \leq 24$.

5. NEW DIRECTION: VECTOR BUNDLES ON CURVES

For G a simple, simply connected complex linear algebraic group, and C a stable curve of arithmetic genus $g \geq 2$, $\text{Bun}_G(C)$ is the stack parameterizing principal G -bundles on C . To any representation $G \rightarrow \text{GL}(V)$, there corresponds a distinguished line bundle $\mathcal{D}(V)$ on $\text{Bun}_G(C)$, the determinant of cohomology line bundle.

In [BG16, Theorem 1.1], we show, when G is $\text{SL}(r)$, and for the standard representation $\text{SL}(r) \rightarrow \text{GL}(V)$, one has that the ring \mathcal{A}_\bullet^C

$$\bigoplus_{m \in \mathbb{Z}_{\geq 0}} \text{H}^0(\text{Bun}_{\text{SL}(r)}(C), \mathcal{D}(V)^m)$$

is finitely generated.

The stacks $\text{Bun}_{\text{SL}(r)}(C)$ are not proper, and so one does not expect that the constituent vector spaces $\text{H}^0(\text{Bun}_{\text{SL}(r)}(C), \mathcal{D}^m)$ to be finite dimensional for any m , much less that their infinite sum would be so. The result is well known for smooth curves C , and in this case: $\text{Proj}(\mathcal{A}_\bullet^C) \cong \text{SU}_C(r)$, where $\text{SU}_C(r)$ is the moduli space parameterizing semistable vector bundles of rank r with trivial determinant on C [BL94, Fal94]. These form a flat family over \mathcal{M}_g , and it is natural to ask if one can extend it to a family over Deligne and Mumford's compactification $\overline{\mathcal{M}}_g$ (This problem was discussed in Section 1: Using conformal blocks, we extend the family).

REFERENCES

- [Ale02] Valery Alexeev, *Complete moduli in the presence of semiabelian group action*, Ann. of Math. (2) **155** (2002), no. 3, 611–708. [↑1](#)
- [AGS14] Valery Alexeev, Angela Gibney, and David Swinarski, *Higher-level \mathfrak{sl}_2 conformal blocks divisors on $\overline{M}_{0,n}$* , Proc. Edinb. Math. Soc. (2) **57** (2014), no. 1, 7–30. [↑2](#)
- [AFSvdW16] Jarod Alper, Maksym Fedorchuk, David Smyth, and Fred van der Wyck, *Second flip in the Hassett-Keel program: A local description*, Compositio Mathematica, to appear (2016), 37. [↑1](#)
- [AFS16a] Jarod Alper, Maksym Fedorchuk, and David Smyth, *Second flip in the Hassett-Keel program: Existence of good moduli spaces*, Compositio Mathematica, to appear (2016), 27. [↑1](#)
- [AFS16b] ———, *Second flip in the Hassett-Keel program: Projectivity*, International Mathematics Research Notices, to appear (2016), 33. [↑1](#)
- [AGSS12] Maxim Arap, Angela Gibney, James Stankewicz, and David Swinarski, *sl_n level 1 conformal blocks divisors on $\overline{M}_{0,n}$* , Int. Math. Res. Not. IMRN **7** (2012), 1634–1680. [↑](#)
- [BL94] Arnaud Beauville and Yves Laszlo, *Conformal blocks and generalized theta functions*, Comm. Math. Phys. **164** (1994), no. 2, 385–419. [↑2, 3](#)
- [BG16] Prakash Belkale and Angela Gibney, *On finite generation of the section ring of the determinant of cohomology line bundle* (2016). on the arxiv, submitted. [↑2, 3](#)
- [BGM16] P. Belkale, A. Gibney, and S. Mukhopadhyay, *Nonvanishing of conformal blocks divisors on $\overline{M}_{0,n}$* , Transform. Groups **21** (2016), no. 2, 329–353. [↑2](#)
- [BGM15] Prakash Belkale, Angela Gibney, and Swarnava Mukhopadhyay, *Vanishing and identities of conformal blocks divisors*, Algebr. Geom. **2** (2015), no. 1, 62–90. [↑2](#)
- [BGK16] Prakash Belkale, Angela Gibney, and Anna Kazanova, *Scaling of conformal blocks and generalized theta functions over $\overline{M}_{g,n}$* , Mathematische Zeitschrift <http://link.springer.com/article/10.1007/s00209-016-1682-1> (2016), 1–27. [↑2](#)
- [BCHM10] Caucher Birkar, Paolo Cascini, Christopher Hacon, and James McKernan, *Existence of minimal models for varieties of log general type*, J. Amer. Math. Soc. **23** (2010), 405–468. [↑1](#)
- [CGK09] L. Chen, A. Gibney, and D. Krashen, *Pointed trees of projective spaces*, J. Algebraic Geom. **18** (2009), no. 3, 477–509. [↑1, 3](#)
- [DM69] P. Deligne and D. Mumford, *The irreducibility of the space of curves of given genus*, Inst. Hautes Études Sci. Publ. Math. **36** (1969), 75–109. [↑1](#)
- [Fak12] Najmuddin Fakhruddin, *Chern classes of conformal blocks*, Compact moduli spaces and vector bundles, Contemp. Math., vol. 564, Amer. Math. Soc., Providence, RI, 2012, pp. 145–176. [↑1, 2](#)
- [Fal94] Gerd Faltings, *A proof for the Verlinde formula*, J. Algebraic Geom. **3** (1994), no. 2, 347–374. [↑2, 3](#)
- [FG03] Gavril Farkas and Angela Gibney, *The Mori cones of moduli spaces of pointed curves of small genus*, Trans. Amer. Math. Soc. **355** (2003), no. 3, 1183–1199 (electronic). [↑3](#)
- [Fed11] Maksym Fedorchuk, *Cyclic Covering Morphisms on $\overline{M}_{0,n}$* (2011). [↑2](#)
- [Gia13] Noah Giansiracusa, *Conformal blocks and rational normal curves*, J. Algebraic Geom. **22** (2013), no. 4, 773–793. [↑2](#)
- [GG12] Noah Giansiracusa and Angela Gibney, *The cone of type A, level 1, conformal blocks divisors*, Adv. Math. **231** (2012), no. 2, 798–814. [↑2](#)
- [GJM13] Noah Giansiracusa, David Jensen, and Han-Bom Moon, *GIT compactifications of $M_{0,n}$ and flips*, Adv. Math. **248** (2013), 242–278. [↑2](#)
- [GKM02] Angela Gibney, Sean Keel, and Ian Morrison, *Towards the ample cone of $\overline{M}_{g,n}$* , J. Amer. Math. Soc. **15** (2002), no. 2, 273–294 (electronic). [↑1, 3](#)
- [GM10] Angela Gibney and Diane Maclagan, *Equations for Chow and Hilbert quotients*, Algebra Number Theory **4** (2010), no. 7, 855–885. [↑3](#)
- [GJMS13] Angela Gibney, David Jensen, Han-Bom Moon, and David Swinarski, *Veronese quotient models of $\overline{M}_{0,n}$ and conformal blocks*, Michigan Math. J. **62** (2013), no. 4, 721–751. [↑2](#)
- [Gib09] Angela Gibney, *Numerical criteria for divisors on \overline{M}_g to be ample*, Compos. Math. **145** (2009), no. 5, 1227–1248. [↑3](#)
- [GM16] Angela Gibney and Swarnava Mukhopadhyay, *On higher Chern classes of vector bundles of conformal blocks* (2016). on the arxiv, and submitted. [↑2](#)
- [GM12] Angela Gibney and Diane Maclagan, *Lower and upper bounds for nef cones*, Int. Math. Res. Not. IMRN **14** (2012), 3224–3255. [↑3](#)
- [Gib12] Angela Gibney, *On extensions of the Torelli map*, Geometry and arithmetic, EMS Ser. Congr. Rep., Eur. Math. Soc., Zürich, 2012, pp. 125–136. [↑2](#)

- [HKT06] Paul Hacking, Sean Keel, and Jenia Tevelev, *Compactification of the moduli space of hyperplane arrangements*, J. Algebraic Geom. **15** (2006), no. 4, 657–680. [↑1](#)
- [HKT09] ———, *Stable pair, tropical, and log canonical compactifications of moduli spaces of del Pezzo surfaces*, Invent. Math. **178** (2009), no. 1, 173–227. [↑1](#)
- [HH13] Brendan Hassett and Donghoon Hyeon, *Log minimal model program for the moduli space of stable curves: the first flip*, Ann. of Math. (2) **177** (2013), no. 3, 911–968. [↑1](#)
- [HH09] ———, *Log canonical models for the moduli space of curves: the first divisorial contraction*, Trans. Amer. Math. Soc. **361** (2009), no. 8, 4471–4489. [↑1](#)
- [HM06] Christopher D. Hacon and James McKernan, *Boundedness of pluricanonical maps of varieties of general type*, Invent. Math. **166** (2006), no. 1, 1–25. [↑1](#)
- [Hob15] Natalie Hobson, *Quantum Koszka and the rank one problem for $\overline{M}_{0,n}$* (2015). [↑2](#)
- [Kap93a] M. M. Kapranov, *Veronese curves and Grothendieck-Knudsen moduli space $\overline{M}_{0,n}$* , J. Algebraic Geom. **2** (1993), no. 2, 239–262. [↑2](#)
- [Kap93b] ———, *Chow quotients of Grassmannians. I*, I. M. Gelfand Seminar, Adv. Soviet Math., vol. 16, Amer. Math. Soc., Providence, RI, 1993, pp. 29–110. [↑2](#)
- [Kaz14] Anna Kazanova, *On S_n invariant conformal blocks vector bundles of rank one on $\overline{M}_{0,n}$* (2014). arXiv:1404.5845v1 [math.AG]. [↑2](#)
- [KM13] Seán Keel and James McKernan, *Contractible extremal rays on $\overline{M}_{0,n}$* , Handbook of moduli. Vol. II, Adv. Lect. Math. (ALM), vol. 25, Int. Press, Somerville, MA, 2013, pp. 115–130. [↑3](#)
- [KNR94] Shrawan Kumar, M. S. Narasimhan, and A. Ramanathan, *Infinite Grassmannians and moduli spaces of G -bundles*, Math. Ann. **300** (1994), no. 1, 41–75. [↑2](#)
- [KSB88] J. Kollár and N. I. Shepherd-Barron, *Threefolds and deformations of surface singularities*, Invent. Math. **91** (1988), no. 2, 299–338. [↑1](#)
- [LS97] Yves Laszlo and Christoph Sorger, *The line bundles on the moduli of parabolic G -bundles over curves and their sections*, Ann. Sci. École Norm. Sup. (4) **30** (1997), no. 4, 499–525 (English, with English and French summaries). [↑2](#)
- [KM76] Finn Faye Knudsen and David Mumford, *The projectivity of the moduli space of stable curves. I. Preliminaries on “det” and “Div”*, Math. Scand. **39** (1976), no. 1, 19–55. [↑1](#)
- [Pau96] Christian Pauly, *Espaces de modules de fibrés paraboliques et blocs conformes*, Duke Math. J. **84** (1996), no. 1, 217–235 (French). [↑2](#)
- [Smy13] David Ishii Smyth, *Towards a classification of modular compactifications of $\mathcal{M}_{g,n}$* , Invent. Math. **192** (2013), no. 2, 459–503. [↑2](#)
- [TK88] Akihiro Tsuchiya and Yukihiro Kanie, *Vertex operators in conformal field theory on \mathbf{P}^1 and monodromy representations of braid group*, Conformal field theory and solvable lattice models (Kyoto, 1986), Adv. Stud. Pure Math., vol. 16, Academic Press, Boston, MA, 1988, pp. 297–372. [↑1](#)
- [TUY89] Akihiro Tsuchiya, Kenji Ueno, and Yasuhiko Yamada, *Conformal field theory on universal family of stable curves with gauge symmetries*, Integrable systems in quantum field theory and statistical mechanics, Adv. Stud. Pure Math., vol. 19, Academic Press, Boston, MA, 1989, pp. 459–566. [↑1](#)