

# ON FACTORIZATION AND VECTOR BUNDLES OF CONFORMAL BLOCKS FROM VERTEX ALGEBRAS

CHIARA DAMIOLINI, ANGELA GIBNEY, AND NICOLA TARASCA

ABSTRACT. Modules over conformal vertex algebras give rise to sheaves of coinvariants and conformal blocks on moduli of stable pointed curves. We show that under certain natural hypotheses, these sheaves satisfy the *factorization* property, a reflection of their inherent combinatorial nature. As an application, we prove they are vector bundles. These provide a generalization of vector bundles defined by integrable modules over affine Lie algebras at a fixed level. Satisfying factorization is essential to a recursive formulation of invariants, like ranks and Chern classes, and to produce new constructions of rational conformal field theories.

By assigning to each marked point on a stable pointed curve a module over a conformal vertex algebra, one can construct a pair of dual vector spaces of coinvariants and conformal blocks. These data give sheaves on the moduli space of stable pointed curves [DGT2]. The main result of this work is that, as was conjectured, e.g., in [Zhu1] and [FBZ], under certain natural hypotheses these sheaves satisfy the *factorization* property (Theorem 1).

That sheaves of coinvariants satisfy factorization and support a projectively flat connection [DGT2] allows one to show they are vector bundles (Theorem 2). Moreover, there are advantages beyond this application.

Factorization enables one to transform fibers of such a bundle to analogous objects defined on curves of smaller genus and with fewer marked points. This leads to recursions and allows one to make inductive arguments. For instance, ranks of bundles defined by integrable modules at a fixed level over affine Lie algebras were computed using factorization (see e.g. [Bea] for an account of the Verlinde formula). An essential ingredient in the formulation of a cohomological field theory, factorization was used in [MOP1, MOP<sup>+</sup>2] to compute the Chern character of these vector bundles on the moduli space  $\overline{\mathcal{M}}_{g,n}$  of stable  $n$ -pointed curves of genus  $g$ , for arbitrary  $g$ , and conclude that the Chern classes lie in the tautological ring.

Conformal field theories can be used to describe classical solutions of the string equations of motion. Rational conformal field theories are constructed with modular functors which arise from vector bundles on the moduli of curves satisfying factorization [MS, Seg1, Seg2]. Modules over Virasoro vertex algebras and vertex algebras associated to affine Lie algebras produce

---

MSC2010. 14H10, 17B69 (primary), 81R10, 81T40, 14D21 (secondary).

*Key words and phrases.* Vertex algebras, conformal blocks and coinvariants, vector bundles on moduli of curves, factorization and sewing.

well-known modular functors giving rational conformal field theories. This work suggests many more.

Let us establish a small amount of notation in order to state our results. Suppose  $(C, P_\bullet)$  is a stable  $n$ -pointed curve, and  $M^1, \dots, M^n$  are modules over a conformal vertex algebra  $V$  (see §1.1 and §1.2). Important to our work is the correspondence between isomorphism classes of irreducible  $V$ -modules and finite-dimensional irreducible  $A(V)$ -modules [Zhu2], where  $A(V)$  is Zhu's associative algebra, described in §1.6.

The vector space of coinvariants (§4.2) is the largest quotient of the tensor product  $\otimes_{i=1}^n M^i$  by the action of a Lie algebra determined by  $V$  and  $(C, P_\bullet)$ . Two Lie algebras have been used to define coinvariants: the chiral Lie algebra  $\mathcal{L}_{C \setminus P_\bullet}(V)$  (§3.1) and Zhu's Lie algebra  $\mathfrak{g}_{C \setminus P_\bullet}(V)$  (§A.1).

Zhu's Lie algebra is defined when the vertex algebra  $V$  is quasi-primary generated, with lowest degree space of dimension one, for either smooth curves [Zhu1, AN], or for rational stable pointed curves [NT]. To show that  $\mathfrak{g}_{C \setminus P_\bullet}(V)$  is a Lie algebra, Zhu uses that any *fixed* smooth algebraic curve admits an atlas such that all transition functions are Möbius transformations. Families of stable rational curves also satisfy this property, a fact used by Nagatomo and Tsuchiya [NT] to extend Zhu's construction to moduli of stable rational curves. Transition functions between charts on families of curves of higher genus are more complicated, and for an arbitrary vertex algebra,  $\mathfrak{g}_{C \setminus P_\bullet}(V)$  is not well-defined.

Here we work with  $\mathcal{L}_{C \setminus P_\bullet}(V)$ , available for any conformal vertex algebra  $V$ , not necessarily quasi-primary generated. Based on work of Beilinson, Feigin, and Mazur for Virasoro algebras [FBZ], the chiral Lie algebra was defined for smooth pointed curves by Frenkel and Ben-Zvi [FBZ, §19.4.14], and shown in [FBZ] to coincide with that studied by Beilinson and Drinfeld in [BD]. In [DGT2], we extend the definition of  $\mathcal{L}_{C \setminus P_\bullet}(V)$  to nodal curves. In §3.3 we give a more detailed description to fit the purpose of this paper.

Coinvariants  $\mathbb{V}(V; M^\bullet)_{(C, P_\bullet)}$  from the chiral Lie algebra are isomorphic to those given by  $\mathfrak{g}_{C \setminus P_\bullet}(V)$  when Zhu's Lie algebra is defined, and the two can be compared (Proposition A.2.1).

Suppose  $C$  has one node  $Q$ , let  $\tilde{C} \rightarrow C$  be the normalization, let  $Q_+, Q_- \in \tilde{C}$  be the preimages of  $Q$ , and set  $Q_\bullet = (Q_+, Q_-)$ . For  $\mathscr{W}$  the set of simple  $V$ -modules and  $W \in \mathscr{W}$ , let  $W'$  be its contragredient module (§1.8).

**Theorem 1** (Factorization). *Let  $V = \oplus_{i \geq 0} V_i$  be a rational and  $C_2$ -cofinite conformal vertex algebra with  $V_0 \cong \mathbb{C}$ . One has*

$$\mathbb{V}(V; M^\bullet)_{(C, P_\bullet)} \cong \bigoplus_{W \in \mathscr{W}} \mathbb{V}(V; M^\bullet \otimes W \otimes W')_{(\tilde{C}, P_\bullet \sqcup Q_\bullet)}.$$

It may be that  $\tilde{C} = C_+ \sqcup C_-$  is disconnected, with  $Q_\pm \in C_\pm$ . Then:

$$\mathbb{V}(V; M^\bullet \otimes W \otimes W')_{(\tilde{C}, P_\bullet \sqcup Q_\bullet)} \cong \mathbb{V}(V; M_+^\bullet \otimes W)_{X_+} \otimes \mathbb{V}(V; M_-^\bullet \otimes W')_{X_-}$$

where  $X_\pm = (C_\pm, P(C_\pm) \sqcup Q_\pm)$ , and  $M_\pm^\bullet$  are the modules at the  $P_\bullet$  on  $C_\pm$ .

The isomorphism of Theorem 1 is constructed in §7. This result and its refinement Theorem 8.3.1 have been known in special cases for about thirty years [BFM, TUY, NT, Cod]. Theorem 1 generalizes the analogous result of Nagatomo and Tsuchiya [NT] for coinvariants defined using Zhu's Lie algebra for stable curves of genus zero, and part of Huang's work in genus zero and one [Hua1, Hua2, Hua4, Hua5]. A brief history of factorization is given in §0.1.

With the aid of Propositions 3.3.1, 5.1.1, and 6.2.1, the adaptation of the outline of the proof in [NT, §8.6] to coinvariants defined using the chiral Lie algebra and for curves of arbitrary genus is made possible. While the coinvariants considered by Nagatomo and Tsuchiya were defined on curves *with formal coordinates*, our results are not dependent on them.

Proposition 3.3.1 gives an explicit description of the chiral Lie algebra on a nodal curve in terms of elements of the chiral Lie algebra on its normalization. To study coinvariants, a module  $Z$  (defined in §6.2) comes into play in the normalization of the curve, when the attaching points of a node are untethered. To work with  $Z$  we require that  $V$  is rational. This implies that, in addition to being finite,  $A(V)$  is semi-simple (§1.7), giving the decomposition of coinvariants on the nodal curve as in the statement of Theorem 1.

Proposition 6.2.1 is applied to reinterpret the (decomposed) coinvariants on the normalization back to coinvariants on the nodal curve. Key to our argument is that the coinvariants we consider are finite-dimensional for smooth curves (Proposition 5.1.1), and so we may study them via their dual spaces, the spaces of conformal blocks, using correlation functions (Remark 4.2.1).

Finite-dimensionality of coinvariants has been known to be true in special cases, and expected to hold in cases where there are finitely many simple modules (see §0.2). A statement about coinvariants defined from the chiral Lie algebra and smooth curves of arbitrary genus, Proposition 5.1.1 is a natural generalization of work of Abe and Nagatomo [AN] for coinvariants defined from Zhu's Lie algebra and smooth curves with formal coordinates. As Abe and Nagatomo, for this result we assume  $V$  is  $C_2$ -cofinite, that is, a certain subspace of  $V$  has finite codimension. This property also implies that  $V$  has finitely many simple modules.

The hypotheses for Theorem 1 can be stated differently to compare with the hypotheses assumed by Nagatomo and Tsuchiya or by Huang (see §0.1 and Remark 1.7.1).

As an application of Theorem 1, we show:

**Theorem 2.** *Let  $V = \bigoplus_{i \geq 0} V_i$  be a rational,  $C_2$ -cofinite, conformal vertex algebra with  $V_0 \cong \mathbb{C}$ , and  $M^\bullet$  an  $n$ -tuple of finitely-generated  $V$ -modules. Then  $\mathbb{V}(V; M^\bullet)$  is a vector bundle of finite rank on  $\overline{\mathcal{M}}_{g,n}$ .*

The proof, presented in §8, follows the path set by [TUY] and [NT]. The result on the interior of  $\overline{\mathcal{M}}_{g,n}$ , the moduli space  $\mathcal{M}_{g,n}$  of smooth  $n$ -pointed curves of genus  $g$ , follows from finite-dimensionality of coinvariants (Proposition 5.1.1) and that the sheaf supports a projectively flat connection [DGT2].

Two ingredients are needed to give Theorem 2 on the whole space  $\overline{\mathcal{M}}_{g,n}$ : Theorem 8.2.1, a more general result on finite-dimensionality of coinvariants, and the *Sewing Theorem* (Theorem 8.3.1), a refined version of factorization. Each of these involve evaluation of the sheaf of coinvariants on a family formed by infinitesimally smoothing a nodal curve (such families are described in §8.1). The proof of the Sewing Theorem (§8), generalizing [NT, Thm 8.4.5] in genus zero, and (part of) [Hua3] in genus zero and one, relies on Theorem 1 and a sewing procedure originally found in [TUY, §6.2]. We can further interpret the Sewing Theorem as a decomposition induced by a differential operator (Remark 8.3.3) which naturally arises from the twisted logarithmic  $\mathcal{D}$ -module structure of sheaves of coinvariants [DGT2, §7].

Examples of vertex algebras satisfying the hypotheses of Theorems 1 and 2 are given in §9. We compute Chern classes of such vector bundles in [DGT1].

**0.1. History of factorization.** Beilinson, Feigin, and Mazur [BFM] showed that factorization holds for sheaves of coinvariants defined by modules over Virasoro algebras. Tsuchiya and Kanie used integrable modules at a fixed level over affine Lie algebras to form such sheaves on moduli of smooth pointed rational curves with coordinates [TK]. Generalized by Tsuchiya, Ueno, and Yamada to moduli of stable pointed coordinatized curves of arbitrary genus, these sheaves were shown to be vector bundles and to satisfy a number of good properties including factorization [TUY]. Tsuchimoto [Tsu] proved the bundles are independent of coordinates and descend to  $\overline{\mathcal{M}}_{g,n}$ . Our arguments follow [NT] in the genus zero case after our study of the chiral Lie algebra allows one to replace Zhu's Lie algebra in the general case.

Factorization was proved for  $g \in \{0, 1\}$  by Huang [Hua1, Hua2, Hua4, Hua5]. His approach was to prove the operator product expansion and the modular invariance of intertwining operators, the two conjectures that Moore and Seiberg formulated and used to derive their polynomial equations [MS]. To construct genus zero and genus one chiral rational conformal field theories, Huang assumes that (i)  $V = \bigoplus_{i \geq 0} V_i$  with  $V_0 \cong \mathbb{C}$ ; (ii) Every  $\mathbb{N}$ -gradable weak  $V$ -module is completely reducible; and (iii)  $V$  is  $C_2$ -cofinite. Our assumptions are (1)  $V = \bigoplus_{i \geq 0} V_i$  with  $V_0 \cong \mathbb{C}$ ; (2)  $V$  is rational; and (3)  $V$  is  $C_2$ -cofinite. Given (i) = (1), conditions (ii) and (iii) of Huang are equivalent to our conditions (2) and (3) (see §1.7 for more details). Huang shows that if one assumes in addition (iv)  $V \cong V'$ , then the modular tensor categories he constructs are *rigid* and *nondegenerate*, and he proves that the Verlinde formula holds.

Codogni in [Cod] proves factorization in case our hypotheses hold, with the additional assumptions (iv)  $V \cong V'$  and (v)  $V$  has no nontrivial modules. Like Nagatomo and Tsuchiya, Codogni works with coinvariants defined on the moduli space of curves with formal coordinates.

We note that the additional assumption (iv) gives that  $V$  is quasi-primary generated and in particular, coinvariants defined from Zhu's Lie algebra and the chiral Lie algebra agree (see §A).

Here we do not assume condition (iv) nor condition (v). This for instance, allows one to apply our results to coinvariants defined from modules over lattice vertex algebras, certain simple  $W$ -algebras, and vertex algebras constructed as orbifolds, commutants, and tensor products, on curves of arbitrary genus, thus providing a large class of new constructions of vector bundles on moduli spaces of stable curves (see §9 for more details).

**0.2. Finite-dimensionality of coinvariants: speculation and prior results.** In [FBZ, page 3 and §5.5.4] the authors single out rational vertex algebras as good candidates for defining finite-dimensional coinvariants (and hence leading to finite-rank vector bundles). Interestingly, it seems that rationality and  $C_2$ -cofiniteness were believed to be equivalent [DLM2, ABD] for vertex algebras at that time (all known cases of vertex algebras having either one of the properties, in fact had both). We now know this is incorrect, as Abe has given a  $C_2$ -cofinite, non-rational conformal vertex algebra [Abe].

Finite-dimensionality of coinvariants has previously been proved in the following special cases: (1) for integrable modules at a fixed level over affine Lie algebras by Tsuchiya, Ueno, and Yamada [TUY, Thm 4.2.4]; (2) modules over  $C_2$ -cofinite Virasoro vertex algebras by Beilinson, Feigin, and Mazur [BFM, §7]; (3) curves of genus zero and modules over  $C_2$ -cofinite conformal vertex algebras  $V = \bigoplus_{i \in \mathbb{N}} V_i$  such that  $V_0 \cong \mathbb{C}$ , by Nagatomo and Tsuchiya [NT, Thm 6.2.1]; (4) smooth curves of positive genus and modules over a quasi-primary generated,  $C_2$ -cofinite conformal vertex algebras  $V = \bigoplus_{i \in \mathbb{N}} V_i$  such that  $V_0 \cong \mathbb{C}$  by Abe and Nagatomo [AN, Thm 4.7].

## 1. BACKGROUND

**1.1. Conformal vertex algebras.** A *conformal vertex algebra* is a 4-tuple  $(V, \mathbf{1}^V, \omega, Y(\cdot, z))$ , throughout simply denoted  $V$  for short, such that:

- (i)  $V = \bigoplus_{i \in \mathbb{Z}_{\geq 0}} V_i$  is a vector space over  $\mathbb{C}$  with  $\dim V_i < \infty$ ;
- (ii)  $\mathbf{1}^V \in V_0$  (the *vacuum vector*), and  $\omega \in V_2$  (the *conformal vector*);
- (iii)  $Y(\cdot, z): V \rightarrow \text{End}(V) \llbracket z, z^{-1} \rrbracket$  is a linear function assigning to every element  $A \in V$  the *vertex operator*  $Y(A, z) := \sum_{i \in \mathbb{Z}} A_{(i)} z^{-i-1}$ .

The datum  $(V, \mathbf{1}^V, \omega, Y(\cdot, z))$  must satisfy the following axioms:

- (a) (*vertex operators are fields*) for all  $A, B \in V$ ,  $A_{(i)}B = 0$ , for  $i \gg 0$ ;
- (b) (*vertex operators of the vacuum*)  $Y(\mathbf{1}^V, z) = \text{id}_V$ :

$$\mathbf{1}_{(-1)}^V = \text{id}_V \quad \text{and} \quad \mathbf{1}_{(i)}^V = 0, \quad \text{for } i \neq -1,$$

and for all  $A \in V$ ,  $Y(A, z)\mathbf{1}^V \in A + zV \llbracket z \rrbracket$ :

$$A_{(-1)}\mathbf{1}^V = A \quad \text{and} \quad A_{(i)}\mathbf{1}^V = 0, \quad \text{for } i \geq 0;$$

- (c) (*weak commutativity*) for all  $A, B \in V$ , there exists an  $N \in \mathbb{Z}_{\geq 0}$  such that

$$(z-w)^N [Y(A, z), Y(B, w)] = 0 \quad \text{in } \text{End}(V) \llbracket z^{\pm 1}, w^{\pm 1} \rrbracket;$$

(d) (*conformal structure*) for  $Y(\omega, z) = \sum_{i \in \mathbb{Z}} \omega_{(i)} z^{-i-1}$ ,

$$[\omega_{(p+1)}, \omega_{(q+1)}] = (p - q) \omega_{(p+q+1)} + \frac{c}{12} \delta_{p+q,0} (p^3 - p) \text{id}_V.$$

Here  $c \in \mathbb{C}$  is the *central charge* of  $V$ . Moreover:

$$\omega_{(1)}|_{V_i} = i \cdot \text{id}_V, \quad \text{for all } i, \quad \text{and} \quad Y(\omega_{(0)}A, z) = \partial_z Y(A, z).$$

1.1.1. *Action of Virasoro.* The conformal structure encodes an action of the Virasoro (Lie) algebra  $\text{Vir}$  on  $V$ . To explain, we recall the definition.

The *Witt (Lie) algebra*  $\text{Der } \mathcal{K}$  represents the functor which assigns to a  $\mathbb{C}$ -algebra  $R$  the Lie algebra  $\text{Der } \mathcal{K}(R) := R((z))\partial_z$  generated over  $R$  by the derivations  $L_p := -z^{p+1}\partial_z$ , for  $p \in \mathbb{Z}$ , with relations  $[L_p, L_q] = (p - q)L_{p+q}$ . The *Virasoro (Lie) algebra*  $\text{Vir}$  represents the functor which assigns to  $R$  the Lie algebra generated over  $R$  by a formal vector  $K$  and the elements  $L_p$ , for  $p \in \mathbb{Z}$ , with Lie bracket given by

$$[K, L_p] = 0, \quad [L_p, L_q] = (p - q)L_{p+q} + \frac{K}{12}(p^3 - p)\delta_{p+q,0}.$$

A representation of  $\text{Vir}$  has *central charge*  $c \in \mathbb{C}$  if  $K \in \text{Vir}$  acts as  $c \cdot \text{id}$ . By making the identification  $L_p = \omega_{(p+1)} \in \text{End}(V)$ , Axiom (d) above implies that  $\text{Vir}$  acts on  $V$  with central charge  $c$ .

1.1.2. *Degree of  $A_{(i)}$ .* As a consequence of the axioms, one has  $A_{(i)}V_k \subseteq V_{k+d-i-1}$  for homogeneous  $A \in V_d$  (see e.g., [Zhu2]). We will then say that the degree of the operator  $A_{(i)}$  is

$$(1) \quad \deg A_{(i)} := \deg(A) - i - 1, \quad \text{for homogeneous } A \text{ in } V.$$

Axiom (d) implies that  $L_0$  acts as a *degree* operator on  $V$ , and  $L_{-1}$ , called the *translation* operator, is determined by  $L_{-1}A = A_{(-2)}\mathbf{1}^V$ , for  $A \in V$ .

1.2. **Modules of conformal vertex algebras.** Let  $(V, \mathbf{1}^V, \omega, Y(\cdot, z))$  be a conformal vertex algebra. A *conformal  $V$ -module*, or simply a  $V$ -module, is a pair  $(M, Y^M(\cdot, z))$ , where:

- (i)  $M = \bigoplus_{i \geq 0} M_i$  is a vector space over  $\mathbb{C}$  with  $\dim M_i < \infty$ ;
- (ii)  $Y^M(\cdot, z): V \rightarrow \text{End}(M) \llbracket z, z^{-1} \rrbracket$  is a linear function that assigns to  $A \in V$  an  $\text{End}(M)$ -valued *vertex operator*  $Y^M(A, z) := \sum_{i \in \mathbb{Z}} A_{(i)}^M z^{-i-1}$ .

The pair  $(M, Y^M(\cdot, z))$  must satisfy the following axioms:

- (a) for all  $A \in V$  and  $v \in M$ , one has  $A_{(i)}^M v = 0$ , for  $i \gg 0$ ;
- (b)  $Y^M(\mathbf{1}^V, z) = \text{id}_M$ ;
- (c) for all  $A, B \in V$ , there exists  $N \in \mathbb{Z}_{\geq 0}$  such that for all  $v \in M$

$$(z - w)^N [Y^M(A, z), Y^M(B, w)] v = 0;$$

- (d) for all  $A \in V$ ,  $v \in M$ , there exists  $N \in \mathbb{Z}_{\geq 0}$ , such that for all  $B \in V$ ,

$$(w + z)^N (Y^M(Y(A, w)B, z) - Y^M(A, w + z)Y^M(B, z)) v = 0;$$

(e) For  $Y^M(\omega, z) = \sum_{i \in \mathbb{Z}} \omega_{(i)}^M z^{-i-1}$ , one has

$$[\omega_{(p+1)}^M, \omega_{(q+1)}^M] = (p - q) \omega_{(p+q+1)}^M + \frac{c}{12} \delta_{p+q,0} (p^3 - p) \text{id}_M,$$

where  $c \in \mathbb{C}$  is the central charge of  $V$ . We identify  $\omega_{(p+1)} \in \text{End}(M)$  with an action of  $L_p \in \text{Vir}$  on  $M$ . Moreover,  $L_0$  acts semi-simply on  $M$ , and  $Y^M(L_{-1}A, z) = \partial_z Y^M(A, z)$ .

We note that  $V$  is a module over itself (see [LL, Thm 3.5.4] or [FBZ, §3.2.1]). In the definition, axiom (c) is referred to as *weak commutativity*, and axiom (d) as *weak associativity*. Weak associativity and weak commutativity are known to be equivalent to the *Jacobi identity* (see for instance [DL], [FHL], [LL], [Li1]). Finally, we note that if  $A \in V$  is homogeneous, then following for example [Zhu2],

$$(2) \quad A_{(i)}^M(M_k) \subseteq M_{k+\text{deg}(A)-i-1}.$$

We refer to Appendix B for a brief summary on variations of the notion of  $V$ -modules.

**1.3. The Lie algebra ancillary to  $V$ .** Given a formal variable  $t$ , in this paper we refer to the quotient

$$\mathfrak{L}_t(V) := (V \otimes \mathbb{C}((t))) / \text{Im } \partial,$$

as the *Lie algebra*  $\mathfrak{L}(V) = \mathfrak{L}_t(V)$  *ancillary to  $V$* . Here

$$(3) \quad \partial := L_{-1} \otimes \text{id}_{\mathbb{C}((t))} + \text{id}_V \otimes \partial_t.$$

Denote by  $A_{[i]}$  the projection in  $\mathfrak{L}(V)$  of  $A \otimes t^i \in V \otimes \mathbb{C}((t))$ . Series of type  $\sum_{i \geq i_0} c_i A_{[i]}$ , for  $A \in V$ ,  $c_i \in \mathbb{C}$ , and  $i_0 \in \mathbb{Z}$ , form a spanning set for  $\mathfrak{L}(V)$ . The Lie bracket of  $\mathfrak{L}(V)$  is induced by

$$(4) \quad [A_{[i]}, B_{[j]}] := \sum_{k \geq 0} \binom{i}{k} (A_{(k)} \cdot B)_{[i+j-k]}.$$

The axiom on the vacuum vector  $\mathbf{1}^V$  implies that  $\mathbf{1}_{[-1]}^V$  is central. The Lie algebra  $\mathfrak{L}(V)$  is isomorphic to the current Lie algebra in [NT]. In the following, we will use, as formal variable  $t$ , a formal coordinate at a point  $P$  on a curve, and we will denote  $\mathfrak{L}_P(V) = \mathfrak{L}_t(V)$ . The Lie algebra  $\mathfrak{L}_t(V)$  has a coordinate-free description, discussed in §2.9.

**1.4. The universal enveloping algebra  $\mathcal{U}(V)$ .** For a vertex algebra  $V$ , there is a complete topological associative algebra  $\mathcal{U}(V)$ , defined originally by Frenkel and Zhu [FZ]. We review it here following the presentation in [FBZ]. Consider the universal enveloping algebra  $U(\mathfrak{L}(V))$  of  $\mathfrak{L}(V)$ , and its completion

$$\widetilde{U}(\mathfrak{L}(V)) := \varprojlim_N U(\mathfrak{L}(V))/I_N,$$

where  $I_N$  is the left ideal generated by  $A_{[i]}$ , for  $A \in V$ , and  $i > N$ . The *universal enveloping algebra*  $\mathcal{U}(V)$  of  $V$  is defined as the quotient of  $\widetilde{U}(\mathfrak{L}(V))$  by the two-sided ideal generated by the Fourier coefficients of the series

$$Y[A_{(-1)}B, z] - :Y[A, z] Y[B, z]:, \quad \text{for all } A, B \in V,$$

where  $Y[A, z] = \sum_{i \in \mathbb{Z}} A_{[i]} z^{-i-1}$ , and the normal ordering  $: \cdot :$  is defined as usual (see e.g., [FLM2]).

Interestingly, for an affine vertex algebra  $V = V_\ell(\mathfrak{g})$ , one has that  $\mathcal{U}(V)$  is isomorphic to a completion  $\widetilde{U}_\ell(\widehat{\mathfrak{g}})$  of  $U_\ell(\widehat{\mathfrak{g}})$ , for all levels  $\ell \in \mathbb{C}$ . Here,  $U_\ell(\widehat{\mathfrak{g}})$  is the quotient of  $U(\widehat{\mathfrak{g}})$  by the two-sided ideal generated by  $K - \ell$ , where  $K \in \widehat{\mathfrak{g}}/\mathfrak{g} \otimes \mathbb{C}((t))$ , and  $\widetilde{U}_\ell(\widehat{\mathfrak{g}})$  is defined as

$$\widetilde{U}_\ell(\widehat{\mathfrak{g}}) := \varprojlim_N U_\ell(\widehat{\mathfrak{g}})/U_\ell(\widehat{\mathfrak{g}}) \cdot \mathfrak{g} \otimes t^N \mathbb{C}[[t]]$$

(see e.g., [FBZ, §4.3.2]). The same holds for vertex algebras  $V$  which are induced representations of a Lie algebra  $\mathfrak{g}$ , such as the Heisenberg and Virasoro vertex algebras. In these cases,  $\mathcal{U}(V)$  is isomorphic to a completion of  $U(\mathfrak{g})$  [FBZ, §5.1.8].

**1.5. Action on  $V$ -modules.** Both  $\mathfrak{L}(V)$  and  $\mathcal{U}(V)$  act on any  $V$ -module  $M$  via the Lie algebra homomorphism  $\mathfrak{L}(V) \rightarrow \text{End}(M)$  obtained by mapping  $A_{[i]}$  to the Fourier coefficient  $A_{(i)}$  of the vertex operator  $Y^M(A, z) = \sum_i A_{(i)} z^{-i-1}$ . More generally, the series  $\sum_{i \geq i_0} c_i A_{[i]}$  acts on a  $V$ -module  $M$  via

$$\text{Res}_{z=0} Y^M(A, z) \sum_{i \geq i_0} c_i z^i dz.$$

We note that an  $\mathfrak{L}(V)$ -module need not be a  $V$ -module. On the other hand, there is an equivalence between the categories of  $V$ -modules and smooth  $\mathcal{U}(V)$ -modules (see [FBZ, §5.1.6]). A  $\mathcal{U}(V)$ -module  $M$  is *smooth* if for any  $w \in M$  and  $A \in V$ , one has  $A_{[i]}w = 0$  for  $i \gg 0$ . Below we review how to describe irreducible  $V$ -modules.

**1.6. Correspondence between  $V$ -modules and  $A(V)$ -modules.** A  $V$ -module  $W$  is *irreducible*, or *simple*, if it has no sub-representation other than the trivial representation 0 and  $W$  itself. We review here Zhu's associative algebra  $A(V)$ , and the one-to-one correspondence between isomorphism classes of finite-dimensional irreducible  $A(V)$ -modules and isomorphism classes of irreducible  $V$ -modules [Zhu2].

*Zhu's algebra* is the quotient  $A(V) := V/O(V)$ , where  $O(V)$  is the vector subspace of  $V$  linearly spanned by elements of the form

$$\text{Res}_{z=0} \frac{(1+z)^{\deg A}}{z^2} Y(A, z)B,$$

where  $A$  is homogeneous in  $V$ . The image of an element  $A \in V$  in  $A(V)$  is denoted by  $o(A)$ . The product in  $A(V)$  is defined by

$$o(A) * o(B) = \operatorname{Res}_{z=0} \frac{(1+z)^{\deg A}}{z} Y(A, z)B,$$

for homogeneous  $A$  in  $V$ . Nagatomo and Tsuchiya [NT] consider an isomorphic copy of  $A(V)$ , which they refer to as the *zero-mode algebra*.

Given a  $V$ -module  $W = \bigoplus_{i \geq 0} W_i$ , one has that  $W_0$  is an  $A(V)$ -module [Zhu2, Thm 2.2.2]. The action of  $A(V)$  on  $W_0$  is defined as follows: an element  $o(A) \in A(V)$ , image of a homogeneous element  $A \in V$ , acts on  $W_0$  as the endomorphism  $A_{(\deg A - 1)}$ , a Fourier coefficient of  $Y^W(A, z)$ .

Conversely, one constructs a  $V$ -module from an  $A(V)$ -module in the following way. The Lie algebra  $\mathfrak{L}(V)$  admits a triangular decomposition:

$$(5) \quad \mathfrak{L}(V) = \mathfrak{L}(V)_{<0} \oplus \mathfrak{L}(V)_0 \oplus \mathfrak{L}(V)_{>0},$$

determined by the degree (1), that is,  $\deg(A_{[i]}) := \deg(A) - i - 1$ , for homogeneous  $A \in V$ . From the definition of the Lie bracket (4) of  $\mathfrak{L}(V)$ , one checks that each summand above is a Lie subalgebra of  $\mathfrak{L}(V)$ . This induces a subalgebra  $\mathcal{U}(V)_{\leq 0}$  of  $\mathcal{U}(V)$ .

Given a finite-dimensional  $A(V)$ -module  $E$ , the *generalized Verma  $\mathcal{U}(V)$ -module* is

$$M(E) := \mathcal{U}(V) \otimes_{\mathcal{U}(V)_{\leq 0}} E.$$

To make  $E$  into an  $\mathcal{U}(V)_{\leq 0}$ -module, one lets  $\mathfrak{L}(V)_{<0}$  act trivially on  $E$ , and  $\mathfrak{L}(V)_0$  act by the homomorphism of Lie algebras  $\mathfrak{L}(V)_0 \rightarrow A(V)_{\text{Lie}}$  induced by the identity endomorphism of  $V$  [Li2, Lemma 3.2.1]. For homogeneous  $A \in V_k$ , the image of the element  $A_{[k-1]} \in \mathfrak{L}(V)_0$  in  $A(V)$  is  $o(A)$ . By construction,  $M(E)$  is automatically a  $V$ -module.

Given an *irreducible*  $V$ -module  $W = \bigoplus_{i \geq 0} W_i$ , the space  $W_0$  is a finite-dimensional *irreducible*  $A(V)$ -module; conversely, given a finite-dimensional *irreducible*  $A(V)$ -module  $E$ , there is a unique maximal proper  $V$ -submodule  $N(E)$  of the  $V$ -module  $M(E)$  with  $N(E) \cap E = 0$  such that  $L(E) = M(E)/N(E)$  is an *irreducible*  $V$ -module [Zhu2].

**1.7. Rational vertex algebras.** Following [FZ], a vertex algebra is *rational* if it has only finitely many isomorphism classes of irreducible modules, and every finitely generated module is a direct sum of irreducible ones. When  $V$  is rational, the associative algebra  $A(V)$  is semisimple [Zhu2].

For a rational vertex algebra  $V$ , given a simple module  $E$  over Zhu's algebra  $A(V)$ , the Verma module  $M(E)$  remains simple. In general, Verma modules are not necessarily simple, but they are indecomposable. Hence complete reducibility implies that simple and indecomposable coincide.

**Remark 1.7.1.** Nagatomo and Tsuchiya asked whether an irreducible  $A(V)$ -module  $E$  always gives rise to an *irreducible*  $V$ -module  $M(E) = L(E)$  when  $V$  is  $C_2$ -cofinite and  $A(V)$  is semisimple [NT, Problem, page 439]. They pointed out that this is indeed the case for rational Virasoro vertex algebras

and for irreducible affine vertex algebras  $L_\ell(\mathfrak{g})$  defined by simple Lie algebras  $\mathfrak{g}$  at level  $\ell$  (see e.g., [NT, §A.1] and §9). As remarked above, given a rational conformal vertex algebra, this irreducibility condition always holds for generalized Verma modules, which are indecomposable, since for a rational vertex algebra, simple and indecomposable modules coincide.

**1.8. Contragredient modules.** Essential to the statement of Theorem 1, contragredient modules are reviewed here following [FHL, §5.2].

Let  $V$  be a vertex algebra. Given a  $V$ -module  $(M = \bigoplus_{i \geq 0} M_i, Y^M(-, z))$ , its *contragredient module* is  $(M', Y^{M'}(-, z))$ , where  $M'$  is the graded dual of  $M$ , that is,  $M' := \bigoplus_{i \geq 0} M_i^\vee$ , with  $M_i^\vee := \text{Hom}_{\mathbb{C}}(M_i, \mathbb{C})$ , and

$$Y^{M'}(-, z): V \rightarrow \text{End}(M') \llbracket z, z^{-1} \rrbracket$$

is the unique linear map determined by

$$(6) \quad \langle Y^{M'}(A, z)\psi, m \rangle = \langle \psi, Y^M(e^{zL_1}(-z^{-2})^{L_0} A, z^{-1}) m \rangle$$

for  $A \in V$ ,  $\psi \in M'$ , and  $m \in M$ . Here and throughout,  $\langle \cdot, \cdot \rangle$  is the natural pairing between a vector space and its graded dual.

1.8.1. In view of the correspondence between isomorphism classes of irreducible  $V$ -modules and irreducible  $A(V)$ -modules (§1.5), we explicitly describe the  $A(V)$ -module structure on  $M_0^\vee$ , with  $M$  a  $V$ -module. For this purpose, consider the involution  $\vartheta: \mathfrak{L}(V) \rightarrow \mathfrak{L}(V)$  induced from

$$(7) \quad \vartheta(A_{[j]}) := (-1)^{k-1} \sum_{i \geq 0} \frac{1}{i!} L_1^i A_{[2k-j-i-2]}$$

for a homogeneous element  $A \in V_k$ . Observe that since the operator  $L_1$  has negative degree, the above sum is finite. This involution appeared in [Bor], and it naturally arises from the action of the vertex operators on the contragredient module  $V'$ . Since  $\vartheta$  restricts to an involution of  $\mathfrak{L}(V)_0$  leaving  $O(V)$  invariant, it induces an involution on Zhu's algebra  $A(V)$ . The following statement is a direct consequence of the definition of contragredient modules.

**Lemma 1.8.1.** *i) The image of  $\psi \in M_0^\vee$  under the action of  $\sigma \in \mathfrak{L}(V)$  is the linear functional*

$$\sigma \cdot \psi = -\psi \circ \vartheta(\sigma).$$

*ii) The image of  $\psi \in M_0^\vee$  under the action of  $o(A) \in A(V)$  is*

$$o(A) \cdot \psi = -\psi \circ \vartheta(o(A)).$$

*For homogeneous  $A \in V$  of degree  $k$ , and for  $m \in M_0$ , this is*

$$\langle o(A) \cdot \psi, m \rangle = (-1)^k \left\langle \psi, \sum_{i \geq 0} \frac{1}{i!} (L_1^i A)_{(k-i-1)} m \right\rangle.$$

We refer the reader to [FBZ, §10.4.8] for a geometric realization of the involution  $\vartheta$ . In §3, the involution  $\vartheta$  will be needed to describe chiral Lie algebras on nodal curves.

**1.9. Stable  $k$ -differentials.** Let  $(C, P_\bullet)$  be a stable  $n$ -pointed curve with at least one node, and let  $\omega_C$  be the dualizing sheaf on  $C$ . We review here (*stable*)  $k$ -differentials on  $C$ , that is, sections of  $\omega_C^{\otimes k}$ , for an integer  $k$ . When  $k \geq 1$ , by  $\omega_C^{\otimes -k}$  we mean  $(\omega_C^\vee)^{\otimes k}$ .

Let  $\tilde{C} \rightarrow C$  be the partial normalization of  $C$  at a node  $Q$ , let  $Q_+, Q_- \in \tilde{C}$  be the two preimages of  $Q$ , and set  $Q_\bullet = (Q_+, Q_-)$ . Note that the curve  $\tilde{C}$  may not be connected. Let  $s_+$  and  $s_-$  be formal coordinates at the points  $Q_+$  and  $Q_-$ , respectively. We write  $Q_\pm$  to denote either point, and similarly use  $s_\pm$  to denote either formal coordinate. For a section

$$\mu \in H^0(\tilde{C} \setminus P_\bullet \sqcup Q_\bullet, \omega_{\tilde{C}}^{\otimes k}) =: \tilde{\Gamma},$$

let  $\mu_{Q_\pm} \in \mathbb{C}((s_\pm))(ds_\pm)^k$  be the Laurent series expansion of  $\mu$  at  $Q_\pm$ , that is, the image of  $\mu$  under the restriction morphism

$$H^0(\tilde{C} \setminus P_\bullet \sqcup Q_\bullet, \omega_{\tilde{C}}^{\otimes k}) \rightarrow H^0(D_{Q_\pm}^\times, \omega_{\tilde{C}}^{\otimes k}) \simeq_{s_\pm} \mathbb{C}((s_\pm))(ds_\pm)^k.$$

Here  $D_{Q_\pm}^\times$  is the punctured formal disk about  $Q_\pm$ , that is, the spectrum of the field of fractions of the completed local ring  $\widehat{\mathcal{O}}_{Q_\pm}$ , and  $\simeq_{s_\pm}$  denotes the isomorphism given by fixing the formal coordinate  $s_\pm$  at  $Q_\pm$ .

For a  $k$ -differential  $\mu$ , define the *order*  $\text{ord}_{Q_\pm}(\mu)$  of  $\mu$  at the point  $Q_\pm$  as the highest integer  $m$  such that  $\mu_{Q_\pm} \in s_\pm^m \mathbb{C}[[s_\pm]](ds_\pm)^k$ . For a  $k$ -differential  $\mu$  with  $\text{ord}_{Q_\pm}(\mu) \geq -k$ , the  *$k$ -residue*  $\text{Res}_{Q_\pm}^k(\mu)$  of  $\mu$  at the point  $Q_\pm$  is defined as the coefficient of  $s_\pm^{-k}(ds_\pm)^k$  in  $\mu_{Q_\pm}$ .

The order and the  $k$ -residue at  $Q_\pm$  are independent of the formal coordinate  $s_\pm$  at  $Q_\pm$ . For the definition of the  $k$ -residue without the assumption  $\text{ord}_{Q_\pm}(\mu) \geq -k$ , see e.g., [BCG<sup>+</sup>]; here we only need the case  $\text{ord}_{Q_\pm}(\mu) \geq -k$ .

**Lemma 1.9.1.** *Assume that  $C \setminus P_\bullet$  is affine. For all integers  $k$ , one has*

$$H^0(C \setminus P_\bullet, \omega_C^{\otimes k}) = \left\{ \mu \in \tilde{\Gamma} \mid \begin{array}{l} \text{ord}_{Q_+}(\mu) \geq -k, \text{ord}_{Q_-}(\mu) \geq -k, \\ \text{Res}_{Q_+}^k(\mu) = (-1)^k \text{Res}_{Q_-}^k(\mu) \end{array} \right\}.$$

*Proof.* It is enough to prove the statement for  $k \in \{-1, 0, 1\}$ : indeed, for negative integers  $k$ , sections of  $\omega_C^{\otimes k}$  on the affine  $C \setminus P_\bullet$  are tensor products of sections of  $\omega_C^{-1}$ , and the Laurent series expansions of sections of  $\omega_C^{\otimes k}$  at  $Q_+$  and  $Q_-$  are obtained as tensors of the Laurent series expansions of sections of  $\omega_C^{-1}$  at  $Q_+$  and  $Q_-$ , respectively. Similarly, for positive integers  $k$ .

When  $k = 1$ , the statement is about sections of  $\omega_C$ , and by definition sections of  $\omega_C$  are sections of  $\omega_{\tilde{C}}$  with at most simple poles at  $Q_+$  and  $Q_-$  such that  $\text{Res}_{Q_+}(\mu) + \text{Res}_{Q_-}(\mu) = 0$ . When  $k = 0$ , the statement is about sections of  $\mathcal{O}_C$ , and indeed a regular function on  $C$  is a regular

function  $\mu$  on  $\widetilde{C}$  such that  $\mu(Q_+) = \mu(Q_-)$ . When  $k = -1$ , by definition we have  $\omega_C^{-1} = \mathcal{H}om_{\widehat{\mathcal{O}}_C}(\omega_C, \widehat{\mathcal{O}}_C)$ , and the statement follows from a direct computation using the cases  $k \in \{0, 1\}$ .  $\square$

**1.10. A consequence of Riemann-Roch.** Throughout this paper, we will use a consequence of the Riemann-Roch theorem which we next describe.

Let  $C$  be a smooth curve, possibly disconnected, with two non-empty sets of distinct marked points  $P_\bullet = (P_1, \dots, P_n)$  and  $Q_\bullet = (Q_1, \dots, Q_m)$ . Assume that each irreducible component of  $C$  contains at least one of the marked points  $P_\bullet$ . In particular, this will imply that  $C \setminus P_\bullet$  is affine. Let  $s_i$  be a formal coordinate at the point  $Q_i$ , for each  $i$ . Fix an integer  $k$ . For all integers  $d$  and  $N$ , there exists  $\mu \in H^0(C \setminus P_\bullet \sqcup Q_\bullet, \omega_C^{\otimes k})$  such that its Laurent series expansions at the points  $Q_\bullet$  satisfy:

$$\begin{aligned} \mu_{Q_i} &\equiv s_i^d (ds_i)^k && \in \mathbb{C}((s_i))(ds_i)^k / s_i^N \mathbb{C}[[s_i]](ds_i)^k, && \text{for a fixed } i, \\ \mu_{Q_j} &\equiv 0 && \in \mathbb{C}((s_j))(ds_j)^k / s_j^N \mathbb{C}[[s_j]](ds_j)^k, && \text{for all } j \neq i. \end{aligned}$$

This statement has appeared for instance in [Zhu1].

## 2. SHEAVES OF VERTEX ALGEBRAS ON STABLE CURVES

Here we describe the sheaf of vertex algebras  $\mathcal{V}_C$  on a stable curve  $C$ . The sheaf  $\mathcal{V}_C$  (§2.6) and its flat connection (§2.8) allow one to give a coordinate-free description of the Lie algebra ancillary to  $V$  (§2.9) and to construct the chiral Lie algebra (§3), whose representations give rise to vector spaces of coinvariants (§4). Introduced in [FBZ] for smooth curves, here we emphasize the construction over nodal curves, carried out in [DGT2].

**2.1. The group scheme  $\text{Aut } \mathcal{O}$ .** Consider the functor which assigns to a  $\mathbb{C}$ -algebra  $R$  the Lie group:

$$\text{Aut } \mathcal{O}(R) = \{z \mapsto \rho(z) = a_1 z + a_2 z^2 + \dots \mid a_i \in R, a_1 \text{ a unit}\}$$

of continuous automorphisms of the algebra  $R[[z]]$  preserving the ideal  $zR[[z]]$ . The group law is given by composition of series:  $\rho_1 \cdot \rho_2 := \rho_2 \circ \rho_1$ . This functor is represented by a group scheme, denoted  $\text{Aut } \mathcal{O}$ .

To construct the sheaf of vertex algebras  $\mathcal{V}_C$  on a stable curve  $C$ , we describe below the principal  $(\text{Aut } \mathcal{O})$ -bundle  $\mathcal{A}ut_C \rightarrow C$ , and actions of  $\text{Aut } \mathcal{O}$  on the conformal vertex algebra  $V$  and on  $\mathcal{A}ut_C \times V$ .

**2.2. Coordinatized curves.** Assume first that  $C$  is a smooth curve. Let  $\mathcal{A}ut_C$  be the infinite-dimensional smooth variety whose points consist of pairs  $(P, t)$ , where  $P$  is a point in  $C$ , and  $t$  is a formal coordinate at  $P$  (see [ADCKP]). A formal coordinate  $t$  at  $P$  is an element of the completed local ring  $\widehat{\mathcal{O}}_P$  such that  $t(P) = 0$  and  $t'(P) \neq 0$ . There is a natural forgetful map  $\mathcal{A}ut_C \rightarrow C$ , with fiber at a point  $P \in C$  equal to the set of formal coordinates at  $P$ :

$$\mathcal{A}ut_P = \{t \in \widehat{\mathcal{O}}_P \mid t(P) = 0, t'(P) \neq 0\}.$$

The group scheme  $\text{Aut } \mathcal{O}$  has a right action on the fibers of  $\mathcal{A}ut_C \rightarrow C$  by change of coordinates:

$$\mathcal{A}ut_C \times \text{Aut } \mathcal{O} \rightarrow \mathcal{A}ut_C, \quad ((P, t), \rho) \mapsto (P, t \cdot \rho := \rho(t)).$$

This action is simply transitive, thus  $\mathcal{A}ut_C$  is a principal  $(\text{Aut } \mathcal{O})$ -bundle on  $C$ . The choice of a formal coordinate  $t$  at  $P$  gives rise to the trivialization

$$\text{Aut } \mathcal{O} \xrightarrow{\cong} \mathcal{A}ut_P, \quad \rho \mapsto \rho(t).$$

For a nodal curve  $C$ , let  $\tilde{C} \rightarrow C$  denote the normalization of  $C$ . A principal  $(\text{Aut } \mathcal{O})$ -bundle on  $C$  is equivalent to the datum of a principal  $(\text{Aut } \mathcal{O})$ -bundle on  $\tilde{C}$  together with gluing isomorphism between the fibers over the two preimages of each node in  $C$ . In particular one constructs the principal  $(\text{Aut } \mathcal{O})$ -bundle  $\mathcal{A}ut_C$  on  $C$  from the principal  $(\text{Aut } \mathcal{O})$ -bundle  $\mathcal{A}ut_{\tilde{C}}$  on  $\tilde{C}$  by identifying the fibers at the two preimages  $Q_+$  and  $Q_-$  of each node  $Q$  in  $C$  by the following gluing identity: if  $C$  is given locally near  $Q$  by the equation  $s_+s_- = 0$ , with  $s_{\pm}$  a formal coordinate at  $Q_{\pm}$ , then the gluing isomorphism is given by

$$(8) \quad \mathcal{A}ut_{Q_+} \simeq_{s_+} \text{Aut } \mathcal{O} \xrightarrow{\cong} \text{Aut } \mathcal{O} \simeq_{s_-} \mathcal{A}ut_{Q_-}, \quad \rho(s_+) \mapsto \rho \circ \gamma(s_-),$$

where  $\gamma \in \text{Aut } \mathcal{O}$  is

$$(9) \quad \gamma(z) = \frac{1}{1+z} - 1 = -z + z^2 - z^3 + \dots$$

Note that  $(\gamma \circ \gamma)(z) = z$ , hence (8) determines an involution of  $\text{Aut } \mathcal{O}$ . The isomorphism (8) is induced from the identification  $s_+ = \gamma(s_-)$ . In §2.2.1 we explain how the element  $\gamma$  arises geometrically from a blow-up construction.

2.2.1. The above construction of  $\mathcal{A}ut_C$  can be carried out in families of stable curves. One thus obtains the moduli space  $\widehat{\mathcal{M}}_{g,1}$  of objects of type  $(C, P, t)$ , where  $(C, P)$  is a stable pointed curve of genus  $g$ , and  $t$  is a formal coordinate at  $P$  (for smooth curves, this moduli space of pointed curves with formal coordinates is constructed in [ADCKP]). For a stable curve  $C$  of genus  $g$ ,  $\mathcal{A}ut_C$  is the fiber of the projection  $\widehat{\mathcal{M}}_{g,1} \rightarrow \overline{\mathcal{M}}_g$  over the point  $[C] \in \overline{\mathcal{M}}_g$ :

$$\begin{array}{ccccc} \mathcal{A}ut_P & \longrightarrow & \mathcal{A}ut_C & \longrightarrow & \widehat{\mathcal{M}}_{g,1} \\ \downarrow & & \downarrow & & \downarrow \text{Aut } \mathcal{O} \\ \text{Spec}(\mathbb{C}) & \xrightarrow{[C,P]} & C & \longrightarrow & \overline{\mathcal{M}}_{g,1} \\ & & \downarrow & & \downarrow \\ & & \text{Spec}(\mathbb{C}) & \xrightarrow{[C]} & \overline{\mathcal{M}}_g. \end{array}$$

Over a singular point  $Q \in C$ , the fiber  $\mathcal{A}ut_Q$  can be identified with the space of formal coordinates at the point  $Q$  in  $C'$ , where  $(C', Q)$  is formed by stable reduction at the unstable pointed curve  $(C, Q)$ . Stable reduction is carried out by blowing-up. As a result,  $C'$  consists of the partial normalization

$\widetilde{C}$  of  $C$  at  $Q$  together with a rational exceptional component meeting  $\widetilde{C}$  transversally at the two preimages  $Q_+$  and  $Q_- \in \widetilde{C}$  of the node  $Q$ . Such a rational component contains three special points: the two attaching points and the point labeled  $Q$ . Up to isomorphism, one can identify this component with  $\mathbb{P}^1$ , with the points attached to  $Q_+$  and  $Q_-$  identified with  $0$  and  $\infty \in \mathbb{P}^1$ , respectively, and the point  $Q$  identified with  $1 \in \mathbb{P}^1$ .

Locally around the two attaching points of  $\mathbb{P}^1$ , the curve  $C'$  is given by equations  $s_+s_0 = 0$  and  $s_\infty s_- = 0$ , where  $s_\pm$  is a formal coordinate at  $Q_\pm$ , and  $s_0, s_\infty$  are formal coordinates at  $0, \infty$ , respectively. The formal coordinates  $s_0$  and  $s_\infty$  satisfy  $s_0 s_\infty = 1$ . A pair of such coordinates induce formal coordinates  $s_0 - 1$  and  $s_\infty - 1$  at  $1 \in \mathbb{P}^1$ , with change of variables given by  $s_0 - 1 = \gamma(s_\infty - 1)$ , with  $\gamma$  as in (9). In particular, the change of variables between the two natural formal coordinates  $s_0 - 1$  and  $s_\infty - 1$  at  $1$  coincide with the identification  $s_+ = \gamma(s_-)$  between the two natural formal coordinates  $s_+$  and  $s_-$  given by the gluing isomorphism (8). It follows that one can identify the fiber  $\mathcal{A}ut_Q$  of  $\mathcal{A}ut_C$  over a node  $Q$  with the space of formal coordinates at the point  $1$  in  $\mathbb{P}^1 \subset C'$ .

For further information about  $\widehat{\mathcal{M}}_{g,1}$ , see [FBZ, §6.5] for the case of smooth curves, and [DGT2, §2] for stable curves.

**2.3. Action of  $\text{Aut } \mathcal{O}$  on  $V$ .** The functor which assigns to a  $\mathbb{C}$ -algebra  $R$  the Lie algebra of the Lie group  $\text{Aut } \mathcal{O}(R)$  is

$$\text{Der}_0 \mathcal{O}(R) = \text{Lie}(\text{Aut } \mathcal{O})(R) = R[[z]]z\partial_z.$$

The Lie algebra  $\text{Der}_0 \mathcal{O}(R)$  is generated over  $R$  by the Virasoro elements  $L_p$ , for  $p \geq 0$ , thus  $\text{Der}_0 \mathcal{O}$  is a Lie subalgebra of the Virasoro Lie algebra.

The action of the Virasoro Lie algebra on a conformal vertex algebra  $V$  restricts to an action of  $\text{Der}_0$  on  $V$ . One can integrate this action to get a left action of  $\text{Aut } \mathcal{O}$  on  $V$  defined as the inductive limit of the actions on the subspaces  $V_{\leq k} := \bigoplus_{i \leq k} V_i$ . This follows from the fact that  $L_0$  acts semi-simply with integral eigenvalues, and  $L_p$  acts locally nilpotently for  $p > 0$  [FBZ, §6.3].

Explicitly, to compute the action on  $V$  of an element  $\rho \in \text{Aut } \mathcal{O}$ , one proceeds as follows. The element  $\rho(z)$  can be expressed as

$$\rho(z) = \exp \left( \sum_{i \geq 0} a_i z^{i+1} \partial_z \right) (z)$$

for some  $a_i \in \mathbb{C}$  (see e.g., [FBZ, §6.3.1]). Assuming  $0 \leq \text{Im}(a_0) < 2\pi$ , the coefficients  $a_i$  are uniquely determined. Hence,  $\rho$  acts on  $V$  as  $\exp(\sum_{i \geq 0} -a_i L_i)$ .

As an example and for later use, the element  $\gamma \in \text{Aut } \mathcal{O}$  from (9) can be expressed as

$$(10) \quad \gamma(z) = e^{-z^2 \partial_z} (-1)^{-z \partial_z} (z).$$

This is a special case of the computation in [FBZ, (10.4.3)]. Thus  $\gamma$  acts on  $V$  as  $e^{L_1(-1)^{L_0}}$ . This element allows one to express the gluing isomorphism for the vertex algebra bundle below.

**2.4. Action of  $\text{Aut } \mathcal{O}$  on  $\mathcal{A}ut_C \times V$ .** The group  $\text{Aut } \mathcal{O}$  has a right equivariant action on the trivial bundle  $\mathcal{A}ut_C \times V \rightarrow \mathcal{A}ut_C$  defined by

$$(P, t, A) \cdot \rho = (P, \rho(t), \rho^{-1} \cdot A),$$

for  $\rho \in \text{Aut } \mathcal{O}$  and  $(P, t, A) \in \mathcal{A}ut_C \times V$ .

**2.5. The vertex algebra bundle.** The quotient of  $\mathcal{A}ut_C \times V$  by the action of  $\text{Aut } \mathcal{O}$  descends along the map  $\mathcal{A}ut_C \rightarrow C$  to the *vertex algebra bundle*  $V_C$  on  $C$ :

$$\begin{array}{ccc} \mathcal{A}ut_C \times V & \longrightarrow & \mathcal{A}ut_C \times_{\text{Aut } \mathcal{O}} V =: V_C \\ \downarrow & & \downarrow \\ \mathcal{A}ut_C & \xrightarrow[\pi]{\text{Aut } \mathcal{O}} & C. \end{array}$$

In  $V_C$ , one has identities

$$(11) \quad (P, t, A) = (P, \rho(t), \rho^{-1} \cdot A),$$

for  $\rho \in \text{Aut } \mathcal{O}$  and  $(P, t, A) \in \mathcal{A}ut_C \times V$ .

**2.6. The sheaf of vertex algebras.** The sheaf of sections of  $V_C$  is the sheaf of vertex algebras:

$$\mathcal{V}_C := (V_C \otimes \pi_* \mathcal{O}_{\mathcal{A}ut_C})^{\text{Aut } \mathcal{O}}.$$

This is a locally free sheaf of  $\mathcal{O}_C$ -modules on  $C$ . The stalk of  $\mathcal{V}_C$  at a point  $P \in C$  is isomorphic to  $\mathcal{A}ut_P \times_{\text{Aut } \mathcal{O}} V$ . Given a formal coordinate  $t$  at  $P$ , one has the trivialization

$$\mathcal{A}ut_P \times_{\text{Aut } \mathcal{O}} V \simeq_t V.$$

For a nodal curve  $C$ , the sheaf  $\mathcal{V}_C$  can be described in terms of the sheaf  $\mathcal{V}_{\tilde{C}}$ , where  $\nu: \tilde{C} \rightarrow C$  is the normalization of  $C$ . For each node  $Q$  of  $C$ , let  $Q_+$  and  $Q_-$  be the two preimages of  $Q$ . Locally around  $Q$ , the curve  $C$  is given by the equation  $s_+ s_- = 0$ , with  $s_{\pm}$  a formal coordinate at  $Q_{\pm}$ . The sheaf  $\mathcal{V}_C$  is isomorphic to a subsheaf of  $\nu_* \mathcal{V}_{\tilde{C}}$ . Such a subsheaf is obtained by identifying the stalks at  $Q_+$  and  $Q_-$  via the gluing isomorphism:

$$V \simeq_{s_+} \mathcal{A}ut_{Q_+} \times_{\text{Aut } \mathcal{O}} V \xrightarrow{\cong} \mathcal{A}ut_{Q_-} \times_{\text{Aut } \mathcal{O}} V \simeq_{s_-} V, \quad A \mapsto e^{L_1(-1)^{L_0}} A.$$

The operator  $e^{L_1(-1)^{L_0}}$  defines an involution on  $V$ , and coincides with the action on  $V$  of the element  $\gamma \in \text{Aut } \mathcal{O}$  from (9) and (10). For homogeneous  $A \in V$  of degree  $k$ , one has

$$\gamma \cdot A = e^{L_1(-1)^{L_0}} A = (-1)^k \sum_{i \geq 0} \frac{1}{i!} L_1^i A.$$

The gluing isomorphism for  $\mathcal{V}_C$  is induced from the gluing isomorphism for  $\mathcal{A}ut_C$  in (8). Indeed, the gluing isomorphism corresponds to the identities

$$(P, s_+, A) = (P, \gamma(s_-), A) = (P, s_-, \gamma \cdot A), \quad \text{in } V_C,$$

for  $(P, t, A) \in \mathcal{A}ut_C \times V$ . The identity  $s_+ = \gamma(s_-)$  follows from (8), and the second equality comes from (11) and the identity  $\gamma = \gamma^{-1}$ .

**2.7. The structure of the sheaf  $\mathcal{V}_C$ .** One has that  $\mathcal{V}_C$  is filtered by  $\mathcal{V}_{\leq k}$ , sheaves of sections of the vector bundles of finite rank  $\mathcal{A}ut_C \times_{\text{Aut } \mathcal{O}} V_{\leq k}$ . We emphasize that while the action of  $\text{Aut } \mathcal{O}$  on  $V_{\leq k}$  is well-defined, the action of  $\text{Aut } \mathcal{O}$  on  $V_k$  is so only modulo  $V_{\leq k-1}$ , for each  $k$ . In particular,  $\mathcal{V}_{\leq k}$  is well-defined, but  $\mathcal{A}ut_C \times_{\text{Aut } \mathcal{O}} V_k$  only makes sense as a quotient of  $\mathcal{A}ut_C \times_{\text{Aut } \mathcal{O}} V_{\leq k}$  modulo  $\mathcal{A}ut_C \times_{\text{Aut } \mathcal{O}} V_{\leq k-1}$ . Consider the associated graded sheaf

$$\text{gr}_\bullet \mathcal{V}_C := \bigoplus_{k \geq 0} \text{gr}_k \mathcal{V}_C, \quad \text{where} \quad \text{gr}_k \mathcal{V}_C := \mathcal{V}_{\leq k} / \mathcal{V}_{\leq k-1}.$$

**Lemma 2.7.1.** *One has*

$$\text{gr}_\bullet \mathcal{V}_C \cong \bigoplus_{k \geq 0} \left( \omega_C^{\otimes -k} \right)^{\oplus \dim V_k}.$$

This was proved in [FBZ, §6.5.9] for smooth curves. The argument made there extends to stable curves if one replaces the sheaf of differentials  $\Omega_C^1$  with the dualizing sheaf  $\omega_C$ . We sketch the proof for the reader's convenience.

*Proof.* Consider  $V_k$  as the quotient  $(\text{Aut } \mathcal{O})$ -representation  $V_{\leq k} / V_{\leq k-1}$ , and let  $A \in V_k$  be nonzero. One has  $L_0 \cdot A = kA$  and  $L_p \cdot A = 0$  in  $V_k = V_{\leq k} / V_{\leq k-1}$  for  $p > 0$ . It follows that  $\mathcal{A} := \mathcal{A}ut_C \times_{\text{Aut } \mathcal{O}} A$  is a line sub-bundle of  $\text{gr}_k \mathcal{V}_C$ . We claim that  $\mathcal{A} \cong \omega_C^{\otimes -k}$ , which implies the statement. To prove the claim, it is enough to identify the transition functions of the two line bundles. When  $C$  is smooth, this is done in [FBZ, §6.5.9]. When  $C$  is nodal,  $\mathcal{A}$  is constructed from the line bundle  $\mathcal{A}ut_{\tilde{C}} \times_{\text{Aut } \mathcal{O}} A$  on the normalization  $\tilde{C}$  of  $C$  by identifying the stalks at the preimages of the nodes. It remains to determine the isomorphisms used to identify these. The gluing isomorphism  $A \mapsto e^{L_1}(-1)^{L_0} A$  of  $\mathcal{V}_{\leq k}$  from §2.6 induces the gluing  $A \mapsto (-1)^{L_0} A$  of  $\text{gr}_k \mathcal{V}_C$ . In particular, this is the gluing isomorphism of  $\mathcal{A}$ , which coincides with the gluing of sections of  $\omega_C^{\otimes -k}$  given by the condition on the residues in Lemma 1.9.1. The assertion therefore follows.  $\square$

As a consequence of Lemma 2.7.1, we record here the following statement, which will be used throughout.

**Lemma 2.7.2.** *Let  $(C, P_\bullet)$  be a stable  $n$ -pointed curve such that  $C \setminus P_\bullet$  is affine. One has:*

$$H^0(C \setminus P_\bullet, \mathcal{V}_C) \cong H^0(C \setminus P_\bullet, \text{gr}_\bullet \mathcal{V}_C).$$

*Proof.* We claim that on the affine open set  $C \setminus P_\bullet$ , one has

$$(12) \quad \mathcal{V}_{\leq k} \cong \bigoplus_{i \leq k} \mathcal{V}_{\leq i} / \mathcal{V}_{\leq i-1} = \text{gr}_{\leq k} \mathcal{V}_C.$$

Assuming (12), then one has, for every  $k \in \mathbb{Z}_{\geq 0}$ , an injection

$$\phi_k: \mathrm{gr}_{\leq k} \mathcal{V}_C \hookrightarrow \mathcal{V}_C,$$

altogether defining a map  $\phi: \mathrm{gr}_{\bullet} \mathcal{V}_C \longrightarrow \mathcal{V}_C$ . The map  $\phi$  gives the isomorphism we seek. To see that  $\phi$  is injective, note that any element  $x$  in  $\mathrm{gr}_{\bullet} \mathcal{V}_C$  is in fact a finite sum, and hence  $x$  is in  $\mathrm{gr}_{\leq k} \mathcal{V}$  for some  $k$ . So if  $x$  is mapped to zero by  $\phi$ , then  $x$  is mapped to zero by  $\phi_k$  for some  $k$ . Since all maps  $\phi_k$  are injective,  $x$  is zero. Surjectivity of  $\phi$  follows from the fact that  $\mathcal{V}_C$  is filtered by the  $\mathcal{V}_{\leq k}$ .

We prove (12) by induction on  $k$  with base case  $k = 0$ . Lemma 2.7.1 implies that  $\mathrm{gr}_k \mathcal{V}_C$  is locally free. On affines, locally free sheaves are projective, and hence on the affine open set  $C \setminus P_{\bullet}$ , the following sequence splits:

$$0 \rightarrow \mathcal{V}_{\leq k-1} \rightarrow \mathcal{V}_{\leq k} \rightarrow \mathrm{gr}_k \mathcal{V}_C \rightarrow 0.$$

In particular, on  $C \setminus P_{\bullet}$

$$\mathcal{V}_{\leq k} \cong \mathcal{V}_{\leq k-1} \oplus \mathrm{gr}_k \mathcal{V}_C \cong \mathrm{gr}_{\leq k-1} \mathcal{V}_C \oplus \mathrm{gr}_k \mathcal{V}_C,$$

and (12) holds.  $\square$

**2.8. The flat connection.** The sheaf  $\mathcal{V}_C$  supports a flat connection

$$\nabla: \mathcal{V}_C \rightarrow \mathcal{V}_C \otimes \omega_C$$

defined for smooth curves in [FBZ, §6] and extended to stable curves in [DGT2, §3.2.4]. On a smooth open set  $U$  in  $C$  admitting a global coordinate  $t$  (e.g., an open  $U$  admitting an étale map  $U \rightarrow \mathbb{A}^1$ ), one has a trivialization  $\mathcal{V}|_U \simeq_t V \times U$ . On  $\mathcal{V}|_U$ , the connection  $\nabla$  is given by  $L_{-1} \otimes \mathrm{id}_U + \mathrm{id}_V \otimes \partial_t$  (compare with (3)).

**2.9. A coordinate-free view of the Lie algebra ancillary to  $V$ .** As a first application of the sheaf of vertex algebras, one obtains a coordinate-free version of the Lie algebra ancillary to  $V$ . Namely, for a smooth point  $P$  in an algebraic curve  $C$  and a formal coordinate  $t$  at  $P$ , one has

$$(13) \quad H^0(D_P^{\times}, \mathcal{V}_C \otimes \omega_C / \mathrm{Im} \nabla) \xrightarrow{\simeq_t} \mathfrak{L}_t(V),$$

where, as before,  $D_P^{\times}$  is the punctured formal disk about  $P$  on  $C$ . The map is given as follows: a section of  $\mathcal{V}_C \otimes \omega_C$  on  $D_P^{\times}$  with respect to the  $t$ -trivialization

$$B \otimes \sum_{i \geq i_0} a_i t^i dt \in V \otimes_{\mathbb{C}} \mathbb{C}((t)) \otimes_{\mathbb{C}((t))} \mathbb{C}((t)) dt \simeq_t H^0(D_P^{\times}, \mathcal{V}_C \otimes \omega_C)$$

maps to

$$\mathrm{Res}_{t=0} Y[B, t] \sum_{i \geq i_0} a_i t^i dt \in \mathfrak{L}_t(V),$$

where  $Y[B, t] := \sum_{i \in \mathbb{Z}} B_{[i]} t^{-i-1}$ . Sections in  $\mathrm{Im} \nabla \subset \mathcal{V}_C \otimes \omega_C$  map to zero, hence this defines a linear map from sections of  $\mathcal{V}_C \otimes \omega_C / \mathrm{Im} \nabla$  on  $D_P^{\times}$  to

$\mathfrak{L}_t(V)$ . Moreover, the vector space  $H^0(D_P^\times, \mathcal{V}_C \otimes \omega_C / \text{Im} \nabla)$  has the structure of a Lie algebra such that (13) is indeed an isomorphism of Lie algebras [FBZ, §§19.4.14, 6.6.9].

### 3. THE CHIRAL LIE ALGEBRA

For a stable  $n$ -pointed curve  $(C, P_\bullet)$  and a conformal vertex algebra  $V$ , we define the chiral Lie algebra  $\mathcal{L}_{C \setminus P_\bullet}(V)$  (§3.1), describe the maps to the Lie algebras ancillary to  $V$  at  $P_i$  (§3.2), and give an explicit description for the chiral Lie algebra on a nodal curve in terms of the normalization of the curve (§3.3). We conclude with a consequence of Riemann-Roch (§3.4).

**3.1. Definition of the chiral Lie algebra.** For  $(C, P_\bullet)$  a stable  $n$ -pointed curve and  $V$  a conformal vertex algebra, set

$$\mathcal{L}_{C \setminus P_\bullet}(V) := H^0(C \setminus P_\bullet, \mathcal{V}_C \otimes \omega_C / \text{Im} \nabla).$$

Here  $\mathcal{V}_C$  and its flat connection  $\nabla$  are as in §2.

**3.2. The chiral Lie algebra maps to the Lie algebra ancillary to  $V$ .** For each  $i$ , let  $t_i$  be a formal coordinate at  $P_i$ , let  $D_{P_i}^\times$  be the punctured formal disk about  $P_i$  on  $C$ , and  $\mathfrak{L}_{t_i}(V)$  be the Lie algebra ancillary to  $V$  (§1.3). Consider the linear map  $\varphi = \varphi_{\mathcal{L}}$  obtained as the composition

$$(14) \quad \varphi_{\mathcal{L}}: \mathcal{L}_{C \setminus P_\bullet}(V) \rightarrow \bigoplus_{i=1}^n H^0(D_{P_i}^\times, \mathcal{V}_C \otimes \omega_C / \text{Im} \nabla) \xrightarrow{\cong} \bigoplus_{i=1}^n \mathfrak{L}_{t_i}(V).$$

The first map is canonical and obtained by restricting sections. The second map is the isomorphism of Lie algebras (13) and depends on the formal coordinates  $t_i$ . After [FBZ, §19.4.14], the first map is a homomorphism of Lie algebras, hence so is  $\varphi$ . The map  $\varphi$  thus induces an action of  $\mathcal{L}_{C \setminus P_\bullet}(V)$  on  $\mathfrak{L}(V)^{\oplus n}$ -modules. This will be used in §4. We remark that [FBZ, §19.4.14] gives a stronger statement: for each  $i$ , the map

$$(15) \quad \mathcal{L}_{C \setminus P_\bullet}(V) \rightarrow H^0(D_{P_i}^\times, \mathcal{V}_C \otimes \omega_C / \text{Im} \nabla) \xrightarrow{\cong} \mathfrak{L}_{t_i}(V)$$

obtained by restricting sections and (13) is also a homomorphism of Lie algebras.

**3.3. A close look at the chiral Lie algebra for nodal curves.** Let  $(C, P_\bullet)$  be a stable  $n$ -pointed curve such that  $C \setminus P_\bullet$  is affine. Assume for simplicity that  $C$  has exactly one node, which we denote by  $Q$ . Let  $\tilde{C} \rightarrow C$  be the normalization of  $C$ , let  $Q_+$  and  $Q_-$  be the two preimages of  $Q$ , and set  $Q_\bullet = (Q_+, Q_-)$ . Let  $s_+$  and  $s_-$  be formal coordinates at  $Q_+$  and  $Q_-$ , respectively, such that locally around  $Q$ , the curve  $C$  is given by the equation  $s_+ s_- = 0$ . The chiral Lie algebra for  $(\tilde{C}, P_\bullet \sqcup Q_\bullet)$  is

$$\mathcal{L}_{\tilde{C} \setminus P_\bullet \sqcup Q_\bullet}(V) = H^0(\tilde{C} \setminus P_\bullet \sqcup Q_\bullet, \mathcal{V}_{\tilde{C}} \otimes \omega_{\tilde{C}} / \text{Im} \nabla),$$

and consider the linear map given by restriction:

$$(16) \quad \mathcal{L}_{\tilde{C} \setminus P_\bullet \sqcup Q_\bullet}(V) \rightarrow H^0(D_{Q_+}^\times, \mathcal{V}_{\tilde{C}} \otimes \omega_{\tilde{C}} / \text{Im} \nabla) \xrightarrow{\simeq_{s_+}} \mathfrak{L}_{Q_+}(V).$$

Recall the triangular decomposition of  $\mathfrak{L}_{Q_{\pm}}(V)$  from (5):

$$\mathfrak{L}_{Q_{\pm}}(V) = \mathfrak{L}_{Q_{\pm}}(V)_{<0} \oplus \mathfrak{L}_{Q_{\pm}}(V)_0 \oplus \mathfrak{L}_{Q_{\pm}}(V)_{>0}.$$

Let  $\sigma_{Q_{\pm}} \in \mathfrak{L}_{Q_{\pm}}(V)$  be the image of  $\sigma \in \mathcal{L}_{\tilde{C} \setminus P_{\bullet} \sqcup Q_{\bullet}}(V)$ , and let  $[\sigma_{Q_{\pm}}]_0$  be the image of  $\sigma_{Q_{\pm}}$  under the projection  $\mathfrak{L}_{Q_{\pm}}(V) \rightarrow \mathfrak{L}_{Q_{\pm}}(V)_0$ .

Recall the involution  $\vartheta$  of  $\mathfrak{L}(V)$  in (7). It restricts to an involution on  $\mathfrak{L}(V)_0$  given for homogeneous  $B \in V_k$  by

$$\vartheta(B_{[k-1]}) = (-1)^{k-1} \sum_{i \geq 0} \frac{1}{i!} L_1^i B_{[k-i-1]}.$$

**Proposition 3.3.1.** *For  $C \setminus P_{\bullet}$  affine, one has*

$$\mathcal{L}_{C \setminus P_{\bullet}}(V) = \left\{ \sigma \in \mathcal{L}_{\tilde{C} \setminus P_{\bullet} \sqcup Q_{\bullet}}(V) \left| \begin{array}{l} \sigma_{Q_+}, \sigma_{Q_-} \in \mathfrak{L}(V)_{\leq 0}, \\ \text{and } [\sigma_{Q_-}]_0 = \vartheta([\sigma_{Q_+}]_0) \end{array} \right. \right\}.$$

*Proof.* Since  $C \setminus P_{\bullet}$  is affine, one has

$$\mathcal{L}_{C \setminus P_{\bullet}}(V) = H^0(C \setminus P_{\bullet}, \mathcal{V}_C \otimes \omega_C) / \nabla H^0(C \setminus P_{\bullet}, \mathcal{V}_C).$$

To study elements in  $\mathcal{L}_{C \setminus P_{\bullet}}(V)$ , we can thus describe their lifts in the vector space  $H^0(C \setminus P_{\bullet}, \mathcal{V}_C \otimes \omega_C)$ . By Lemmas 2.7.1 and 2.7.2, one has

$$(17) \quad H^0(C \setminus P_{\bullet}, \mathcal{V}_C \otimes \omega_C) \cong \bigoplus_{k \geq 0} H^0\left(C \setminus P_{\bullet}, \left(\omega_C^{\otimes 1-k}\right)^{\oplus \dim V_k}\right).$$

From the proof of Lemma 2.7.2, this isomorphism is given by the splitting of exact sequences of vector bundles on the affine set  $C \setminus P_{\bullet}$ , one sequence for each degree  $k$ . In the direct sum decomposition of the right-hand side of (17), the transition functions for the sheaf  $\mathcal{V}_C \otimes \omega_C$  are represented by block upper-triangular matrices, with the diagonal blocks giving the transition functions for  $\left(\omega_C^{\otimes 1-k}\right)^{\oplus \dim V_k}$ . This comes from a general property of transition functions for split short exact sequences, see e.g., [Vak, §13.5.A].

*Step 1.* Let  $\tilde{C}$  be the normalization of  $C$ . After Lemma 1.9.1, a section of  $\omega_C^{\otimes 1-k}$  can be described as a section  $\mu$  of  $\omega_{\tilde{C}}^{\otimes 1-k}$  on  $\tilde{C} \setminus P_{\bullet} \sqcup Q_{\bullet}$  such that

$$(18) \quad \text{ord}_{Q_{\pm}}(\mu) \geq k-1 \quad \text{and} \quad \text{Res}_{Q_+}^{1-k}(\mu) = (-1)^{1-k} \text{Res}_{Q_-}^{1-k}(\mu).$$

The above residue condition coincides with the gluing isomorphism of stable  $(1-k)$ -differentials. Under the isomorphism (17), a section of the left-hand side of (17) can be described as a section  $\sigma$  in

$$(19) \quad \bigoplus_{k \geq 0} H^0\left(\tilde{C} \setminus P_{\bullet} \sqcup Q_{\bullet}, \left(\omega_{\tilde{C}}^{\otimes 1-k}\right)^{\oplus \dim V_k}\right)$$

satisfying constraints given by the order and gluing conditions that as we next show, give the assertion. Let us first set some notation. Consider the

composition of the linear maps

(20)

$$\begin{array}{ccc}
\bigoplus_{k \geq 0} H^0 \left( \tilde{C} \setminus P_\bullet \sqcup Q_\bullet, \left( \omega_{\tilde{C}}^{1-k} \right)^{\oplus \dim V_k} \right) & \dashrightarrow & V \otimes_{\mathbb{C}} \mathbb{C}((s_\pm)) ds_\pm \\
\downarrow & & \uparrow \cong \\
\bigoplus_{k \geq 0} H^0 \left( D_{Q_\pm}^\times, \left( \omega_{\tilde{C}}^{1-k} \right)^{\oplus \dim V_k} \right) & \xrightarrow{\cong_{s_\pm}} & \bigoplus_{k \geq 0} \left( \mathbb{C}(ds_\pm)^{-k} \right)^{\oplus \dim V_k} \otimes \mathbb{C}((s_\pm)) ds_\pm
\end{array}$$

where the left vertical map is the restriction of sections, followed by the  $s_\pm$ -trivialization, and then the isomorphism of vector spaces induced by

$$V \cong \bigoplus_{k \geq 0} \left( \mathbb{C}(ds_\pm)^{-k} \right)^{\oplus \dim V_k}.$$

Denote by  $\sigma_{Q_\pm}$  the image of  $\sigma$  via the composition of the maps in (20). The image of  $\sigma_{Q_\pm}$  in  $\mathfrak{L}(V)$  is still denoted by  $\sigma_{Q_\pm}$ .

*Step 2.* We now show how the order conditions from (18) imply  $\sigma_{Q_\pm} \in \mathfrak{L}(V)_{\leq 0}$ . The inequalities  $\text{ord}_{Q_\pm}(\mu) \geq k - 1$  required for sections  $\mu$  of  $\omega_{\tilde{C}}^{\otimes 1-k}$  imply that the image of the section  $\sigma$  via the composition of the maps in (20) lies in

$$(21) \quad \bigoplus_{k \geq 0} V_k \otimes_{\mathbb{C}} s_\pm^{k-1} \mathbb{C}[[s_\pm]] ds_\pm.$$

In other words,  $\sigma$  is required to be in the following subset of (19):

$$(22) \quad \bigoplus_{k \geq 0} H^0 \left( \tilde{C} \setminus P_\bullet, \left( \omega_{\tilde{C}}(Q_+ + Q_-)^{\otimes 1-k} \right)^{\oplus \dim V_k} \right).$$

Recall that the image of an element  $B \otimes \mu \in V \otimes \mathbb{C}((s_\pm)) ds_\pm$  via the projection  $V \otimes \mathbb{C}((s_\pm)) ds_\pm \rightarrow \mathfrak{L}_{Q_\pm}(V)$  is

$$\text{Res}_{s_\pm=0} Y[B, s_\pm] \mu \quad \in \quad \mathfrak{L}_{Q_\pm}(V).$$

This implies that the image of (21) in  $\mathfrak{L}_{Q_\pm}(V)$  lies in  $\mathfrak{L}_{Q_\pm}(V)_{\leq 0}$ , hence one has  $\sigma_{Q_\pm} \in \mathfrak{L}_{Q_\pm}(V)_{\leq 0}$ .

*Step 3.* Finally, we show how the gluing isomorphism identifying elements of  $H^0(C \setminus P_\bullet, \mathcal{V}_C \otimes \omega_C)$  inside (22) implies  $[\sigma_{Q_-}]_0 = \vartheta([\sigma_{Q_+}]_0)$ . For a section  $\sigma$  in (22),  $[\sigma_{Q_\pm}]_0$  denotes the image of  $\sigma$  via the composition of (20) with the projection

$$V \otimes_{\mathbb{C}} \mathbb{C}((s_\pm)) ds_\pm \rightarrow \bigoplus_{k \geq 0} V_k \otimes_{\mathbb{C}} \mathbb{C}s_\pm^{k-1} ds_\pm.$$

The degree  $k$  component of  $[\sigma_{Q_\pm}]_0$  coincides with the  $(1-k)$ -residue at  $Q_\pm$  of the degree  $k$  component of  $\sigma$ . For  $\sigma$  to correspond to a section of the left-hand side of (17), the two elements  $[\sigma_{Q_\pm}]_0$  at the punctured formal disks  $D_{Q_\pm}^\times$  need to satisfy an identity coming from the gluing isomorphism between the stalks at  $Q_\pm$ .

The gluing isomorphism identifying stalks at  $Q_\pm$  of  $\mathcal{V}_{\tilde{C}}$  from §2.6 induces the following gluing isomorphism of stalks at  $Q_\pm$  of (22):

$$\begin{array}{ccc}
 \bigoplus_{k \geq 0} (\mathbb{C}s_+^k ds_+^{-k})^{\oplus \dim V_k} \otimes_{\mathbb{C}} \mathbb{C}s_+^{-1} ds_+ & \xrightarrow{\cong} & \bigoplus_{k \geq 0} (\mathbb{C}s_-^k ds_-^{-k})^{\oplus \dim V_k} \otimes_{\mathbb{C}} \mathbb{C}s_-^{-1} ds_- \\
 \Downarrow \uparrow & & \Downarrow \uparrow \\
 V \otimes_{\mathbb{C}} \mathbb{C}s_+^{-1} ds_+ & & V \otimes_{\mathbb{C}} \mathbb{C}s_-^{-1} ds_- \\
 \Downarrow & & \Downarrow \\
 A \otimes s_+^{-1} ds_+ & \longmapsto & - (e^{L_1(-1)^{L_0}} A) \otimes s_-^{-1} ds_-.
 \end{array}$$

As for the transition functions discussed at the beginning of the proof, in the direct sum decomposition of the stalks, this gluing isomorphism is represented by block upper-triangular matrices with the diagonal blocks giving the gluing isomorphism of the bundles  $(\omega_C^{\otimes 1-k})^{\oplus \dim V_k}$ , namely the residue conditions in (18). The isomorphism between  $V \otimes_{\mathbb{C}} \mathbb{C}s_{\pm}^{-1} ds_{\pm}$  and the stalk of  $\mathcal{L}_{\tilde{C} \setminus P_{\bullet} \sqcup Q_{\bullet}}(V)$  at  $Q_{\pm}$  maps  $L_1^i A \otimes s_{\pm}^{-1} ds_{\pm}$  to  $A_{[k-i-1]}$ , for  $A \in V_k$ . It follows that

$$[\sigma_{Q_+}]_0 = -e^{L_1(-1)^{L_0}} [\sigma_{Q_-}]_0 = \vartheta([\sigma_{Q_-}]_0),$$

hence the statement.  $\square$

**3.4. A consequence of Riemann-Roch for chiral Lie algebras.** In parallel with §1.10, we have the following statement for chiral Lie algebras. Let  $C$  be a smooth curve, possibly disconnected, with two non-empty sets of distinct marked points  $P_{\bullet} = (P_1, \dots, P_n)$  and  $Q_{\bullet} = (Q_1, \dots, Q_m)$ . For each  $i \in \{1, \dots, m\}$ , let  $s_i$  be a formal coordinate at the point  $Q_i$ . For a section  $\sigma \in \mathcal{L}_{C \setminus P_{\bullet} \sqcup Q_{\bullet}}(V)$ , let  $\sigma_{Q_i}$  be the image of  $\sigma$  under the map given by restriction

$$\mathcal{L}_{C \setminus P_{\bullet} \sqcup Q_{\bullet}}(V) \rightarrow H^0(D_{Q_i}^{\times}, \mathcal{V}_C \otimes \omega_C / \text{Im } \nabla) \xrightarrow{\simeq_{s_i}} \mathfrak{L}_{Q_i}(V).$$

For an integer  $N$ , consider

$$\mathfrak{L}_{Q_i}(V, NQ_i) = V \otimes s_i^N \mathbb{C}[[s_i]] / \text{Im } \partial.$$

This is a Lie subalgebra of  $\mathfrak{L}_{Q_i}(V)$ .

**Proposition 3.4.1.** *Assume that  $C \setminus P_{\bullet}$  is affine. Fix a homogeneous element  $E \in V$ , and integers  $d$  and  $N$ . There exists  $\sigma \in \mathcal{L}_{C \setminus P_{\bullet} \sqcup Q_{\bullet}}(V)$  such that:*

$$\begin{array}{lll}
 \sigma_{Q_i} \equiv E_{[d]} & \in & \mathfrak{L}_{Q_i}(V) / \mathfrak{L}_{Q_i}(V, NQ_i), \quad \text{for a fixed } i, \\
 \sigma_{Q_j} \equiv 0 & \in & \mathfrak{L}_{Q_j}(V) / \mathfrak{L}_{Q_j}(V, NQ_j), \quad \text{for all } j \neq i.
 \end{array}$$

*Proof.* Since  $C \setminus P_{\bullet}$  is affine, so is  $C \setminus P_{\bullet} \sqcup Q_{\bullet}$ . As in Proof of Proposition 3.3.1, elements of  $\mathcal{L}_{C \setminus P_{\bullet} \sqcup Q_{\bullet}}(V)$  can be lifted to sections of  $\mathcal{V}_C \otimes \omega_C$  on  $C \setminus P_{\bullet} \sqcup Q_{\bullet}$ , and thus described as sections of  $\bigoplus_{k \geq 0} (\omega_C^{\otimes 1-k})^{\oplus \dim V_k}$  on  $C \setminus P_{\bullet} \sqcup Q_{\bullet}$  via the isomorphism (17). The statement thus follows from the analogous property of sections of tensor products of  $\omega_C$ , discussed in §1.10.  $\square$

## 4. SPACES OF COINVARIANTS

Given a stable pointed curve  $(C, P_\bullet)$  and a conformal vertex algebra  $V$ , we define spaces of coinvariants using representations of the chiral Lie algebra.

**4.1. Representations of the chiral Lie algebra.** We begin by defining the action of the chiral Lie algebra  $\mathcal{L}_{C \setminus P_\bullet}(V)$  on  $M^\bullet := M^1 \otimes \cdots \otimes M^n$ , for  $V$ -modules  $M^1, \dots, M^n$ . For each  $i$ , let  $t_i$  be a formal coordinate at  $P_i$ , and  $\mathfrak{L}_{t_i}(V)$  be the Lie algebra ancillary to  $V$  (§1.3). Each  $\mathfrak{L}_{t_i}(V)$  acts on the  $V$ -module  $M^i$  as in §1.3, and the sum  $\bigoplus_{i=1}^n \mathfrak{L}_{t_i}(V)$  acts diagonally on  $M^\bullet$ . The map (14) thus induces an action of  $\mathcal{L}_{C \setminus P_\bullet}(V)$  on  $M^\bullet$  as follows: for  $\sigma \in \mathcal{L}_{C \setminus P_\bullet}(V)$  and  $A^i \in M^i$ , one has

$$\sigma(A^1 \otimes \cdots \otimes A^n) = \sum_{i=1}^n A^1 \otimes \cdots \otimes \sigma_{P_i}(A^i) \otimes \cdots \otimes A^n,$$

where  $\sigma_{P_i}$  is the restriction of the section  $\sigma$  to the punctured formal disk  $D_{P_i}^\times$  about  $P_i$  on  $C$ .

**4.2. Coinvariants.** When  $C \setminus P_\bullet$  is affine, the *space of coinvariants* at  $(C, P_\bullet, t_\bullet)$  is

$$\mathbb{V}(V; M^\bullet)_{(C, P_\bullet, t_\bullet)} := M_{\mathcal{L}_{C \setminus P_\bullet}(V)}^\bullet = M^\bullet / \mathcal{L}_{C \setminus P_\bullet}(V) \cdot M^\bullet.$$

This is the largest quotient of  $M^\bullet$  on which  $\mathcal{L}_{C \setminus P_\bullet}(V)$  acts trivially. In general, when  $C \setminus P_\bullet$  is not necessarily affine, the *space of coinvariants* at  $(C, P_\bullet, t_\bullet)$  is defined as the direct limit

$$(23) \quad \mathbb{V}(V; M^\bullet)_{(C, P_\bullet, t_\bullet)} := \varinjlim_{(Q_\bullet, s_\bullet)} \mathbb{V}(V; M^\bullet \sqcup (V, \dots, V))_{(C, P_\bullet \sqcup Q_\bullet, t_\bullet \sqcup s_\bullet)}$$

where  $Q_\bullet = (Q_1, \dots, Q_m)$  ranges over the set of stable points of  $C$  such that  $P_\bullet \cap Q_\bullet = \emptyset$  and  $C \setminus P_\bullet \sqcup Q_\bullet$  is affine, and  $s_\bullet = (s_1, \dots, s_m)$ , with  $s_i$  a formal coordinate at  $Q_i$ , for each  $i$ . The above direct limit is well defined thanks to propagation of vacua (this is similar to the case of affine Lie algebras [Fak, Loo], see [DGT2] for more details).

The vector spaces  $\mathbb{V}(V; M^\bullet)_{(C, P_\bullet, t_\bullet)}$  were initially considered for stable pointed curves and modules of Virasoro algebras in [BFM], generalized to conformal vertex algebras and smooth pointed curves in [FBZ, BD], and extended to stable pointed curves in [DGT2]. Since they are *independent of coordinates*, one can define vector spaces of coinvariants, denoted  $\mathbb{V}(V; M^\bullet)_{(C, P_\bullet)}$ , non-canonically isomorphic to  $\mathbb{V}(V; M^\bullet)_{(C, P_\bullet, t_\bullet)}$  for each  $t_\bullet$  (see [DGT2] for more details).

**Remark 4.2.1.** In §5 we show that the vector spaces of coinvariants are finite-dimensional under some natural hypotheses. In particular, one can detect many aspects of the vector spaces of coinvariants by studying their dual spaces, referred to in the literature as vector spaces of *conformal blocks*. Conformal blocks are often studied through their identification with systems of correlation functions [TUY, §2.4], [TK, §5], [NT, §5.3], [FZ, §2.3], [FBZ, §§4.5, 10.3], and [Zhu2, §6]. For instance, systems of correlation functions

were used in the proofs of *propagation of vacua* and the *factorization property* for conformal blocks defined by modules over affine Lie algebras [TUY], in [NT, Thms 5.6.1 and 8.4.3] for conformal blocks in genus zero defined by modules over vertex algebras, and in [Zhu1, Thm 6.1], and [FBZ, §10.3.1] to prove propagation of vacua for conformal blocks in positive genus defined by modules over vertex algebras. In [FZ, Thms 2.3.3 and 2.3.4] correlation functions are used to construct vertex algebras and their representations. While we only nominally use this identification in Remark 6.2.2 and Proposition 6.2.1 in service of the proof of Theorem 1, it is natural to ask whether, as for those associated to vertex algebras induced from affine Lie algebras, the Heisenberg Lie algebra, and the Virasoro algebra [KZ], [MMS], systems of correlation functions are governed by interesting differential equations.

## 5. FINITE-DIMENSIONALITY OF COINVARIANTS

Using coinvariants by the action of Zhu's Lie algebra (§A), Abe and Nagatomo show that spaces of coinvariants at smooth pointed curves of arbitrary genus are finite-dimensional [AN]. To be well-defined, Zhu's Lie algebra requires that the vertex algebra is quasi-primary generated. The chiral Lie algebra provides a generalization of Zhu's Lie algebra beyond this constraint. We show here that the result of [AN] extends to coinvariants by the action of the chiral Lie algebra. Moreover, we further extend the result in [AN] by allowing the following twist of the chiral Lie algebra: given a smooth  $n$ -pointed curve  $(C, P_\bullet)$ , and an effective divisor  $D = \sum_{i=1}^m n_i Q_i$  on  $C$  not supported at  $P_\bullet$ , consider

$$(24) \quad \mathcal{L}_{C \setminus P_\bullet}(V, D) := H^0(C \setminus P_\bullet, \mathcal{V}_C \otimes \omega_C(-D)/\text{Im} \nabla),$$

where  $\text{Im} \nabla$  denotes the restriction of  $\nabla(\mathcal{V}_C)$  to  $\mathcal{V}_C \otimes \omega_C(-D)$ . This is the space of sections in  $\mathcal{L}_{C \setminus P_\bullet}(V)$  vanishing with order at least  $n_i$  at  $Q_i$ , for each  $i$ , and gives a Lie subalgebra of  $\mathcal{L}_{C \setminus P_\bullet}(V)$ . To state the result, we review the definition of  $C_2$ -cofiniteness.

**5.1.  $C_2$ -cofiniteness.** Let  $M$  be a  $V$ -module (e.g.,  $M = V$ ) and consider the following subset of  $M$  for  $k \geq 2$ :

$$C_k(M) := \text{span}_{\mathbb{C}} \{A_{(-k)}m : A \in V, m \in M\}.$$

One says that  $M$  is  $C_k$ -cofinite if  $\dim_{\mathbb{C}} M/C_k(M) < \infty$ . If  $V = \bigoplus_{i \geq 0} V_i$  is a  $C_2$ -cofinite conformal vertex algebra with  $V_0 \simeq \mathbb{C}$ , then any finitely generated  $V$ -module is  $C_k$ -cofinite, for  $k \geq 2$  [Buh]. As explained in [Ara1], the  $C_2$ -cofiniteness has a natural geometric interpretation which generalizes the concept of lisse modules introduced in [BFM] for the Virasoro algebra.

**Proposition 5.1.1.** *Let  $V = \bigoplus_{i \geq 0} V_i$  be a  $C_2$ -cofinite conformal vertex algebra with  $V_0 \cong \mathbb{C}$ . Let  $C$  be a smooth curve with distinct points  $P_1, \dots, P_n$ , and  $D$  an effective divisor on  $C$  not supported at  $P_\bullet$ . For finitely generated  $V$ -modules  $M^1, \dots, M^n$ , the coinvariants  $M_{\mathcal{L}_{C \setminus P_\bullet}(V, D)}^\bullet$  are finite-dimensional.*

*Proof.* Fix formal coordinates  $t_i$  at  $P_i$ , for each  $i$ , and recall the map from (15):  $\mathcal{L}_{C \setminus P_\bullet}(V, D) \rightarrow \mathfrak{L}_{P_i}(V)$ ,  $\sigma \mapsto \sigma_{P_i}$ . Define a filtration on  $\mathcal{L}_{C \setminus P_\bullet}(V, D)$  by

$$\mathcal{F}_k \mathcal{L}_{C \setminus P_\bullet}(V, D) := \left\{ \sigma \in \mathcal{L}_{C \setminus P_\bullet}(V, D) \mid \deg \sigma_{P_i} \leq k, \text{ for all } i \right\}$$

for  $k \in \mathbb{N}$ , giving  $\mathcal{L}_{C \setminus P_\bullet}(V, D)$  the structure of a filtered Lie algebra. Consider also the filtration on  $M^\bullet$  given by

$$\mathcal{F}_k M^\bullet = \bigoplus_{0 \leq d \leq k} M_d^\bullet, \quad \text{where } M_d^\bullet := \sum_{d_1 + \dots + d_n = d} M_{d_1}^1 \otimes \dots \otimes M_{d_n}^n.$$

Since  $\mathcal{F}_k \mathcal{L}_{C \setminus P_\bullet}(V, D) \cdot \mathcal{F}_l M^\bullet \subset \mathcal{F}_{k+l} M^\bullet$ , the  $\mathcal{L}_{C \setminus P_\bullet}(V, D)$ -module  $M^\bullet$  is a filtered  $\mathcal{L}_{C \setminus P_\bullet}(V, D)$ -module. Finally, one has an induced filtration on  $M_{\mathcal{L}_{C \setminus P_\bullet}(V, D)}^\bullet$ :

$$\mathcal{F}_k \left( M_{\mathcal{L}_{C \setminus P_\bullet}(V, D)}^\bullet \right) := \left( \mathcal{F}_k M^\bullet + \mathcal{L}_{C \setminus P_\bullet}(V, D) \cdot M^\bullet \right) / \mathcal{L}_{C \setminus P_\bullet}(V, D) \cdot M^\bullet.$$

*Step 1.* Let  $U$  be a finite-dimensional subspace of  $V$  such that  $V = U \oplus C_2(V)$ . Contrary to [AN], elements of  $U$  are not required to be quasi-primary here. Let  $d_U$  be the maximum of the degree of the homogeneous elements in  $U$ . Similar to [AN, Lemma 4.1], by an application of the Riemann-Roch and the Weierstrass gap theorem, there exists an integer  $N$  such that

$$H^0 \left( C, \omega_C^{\otimes 1-k} (lP_i - D) \right) \neq \emptyset, \quad \text{for all } k \leq d_U, l \geq N, i \in \{1, \dots, n\}.$$

*Step 2.* For a  $V$ -module  $M$  and with  $N$  as in Step 1, define the subset

$$C_N(U, M) = \text{span}_{\mathbb{C}} \left\{ A_{(-l)} m : A \in U, m \in M, l \geq N \right\}.$$

We claim that for each  $i$  the set  $M^1 \otimes \dots \otimes C_N(U, M^i) \otimes \dots \otimes M^n$  is in the kernel of the canonical surjective linear map

$$M^\bullet \xrightarrow{\pi} \text{gr}_\bullet \left( M_{\mathcal{L}_{C \setminus P_\bullet}(V, D)}^\bullet \right) := \bigoplus_{k \geq 0} \mathcal{F}_k \left( M_{\mathcal{L}_{C \setminus P_\bullet}(V, D)}^\bullet \right) / \mathcal{F}_{k-1} \left( M_{\mathcal{L}_{C \setminus P_\bullet}(V, D)}^\bullet \right).$$

For this, it is enough to show that  $\pi \left( m_1 \otimes \dots \otimes A_{(-l)} m_i \otimes \dots \otimes m_n \right) = 0$ , for homogeneous  $A \in U$  of degree  $k$ ,  $m_i \in M_{d_i}^i$ , and  $l \geq N$ . Note that  $C \setminus P_i$  is affine, for all  $i$ . As in the proof of Proposition 3.3.1, elements of  $\mathcal{L}_{C \setminus P_i}(V, D) \subset \mathcal{L}_{C \setminus P_\bullet}(V, D)$  can be lifted to sections of  $\mathcal{V}_C \otimes \omega_C(-D)$  on  $C \setminus P_i$ . By Lemmas 2.7.2 and 2.7.1, the vector space of such sections is isomorphic to the space of sections of

$$(25) \quad \bigoplus_{k \geq 0} V_k \otimes \omega_C^{\otimes 1-k}(-D)$$

on  $C \setminus P_i$ . Following Step 1, there exists a section  $\sigma = A \otimes \mu$  of (25) on  $C \setminus P_i$  such that its image via the map  $\mathcal{L}_{C \setminus P_i}(V, D) \rightarrow \mathfrak{L}_{P_i}(V)$  from (15) is

$$\sigma_{P_i} = A_{[-l]} + \sum_{j > -l} c_j A_{[j]}, \quad \text{for some } c_j \in \mathbb{C}.$$

One has  $A_{[-l]} \cdot M_{d_i}^i \subset M_{d_i+k+l-1}^i$  and  $A_{[j]} \cdot M_{d_i}^i \subset M_{d_i+k+l-2}^i$  for  $j > -l$ . Moreover, since  $\mu$  is holomorphic at a point  $P_j \neq P_i$ , one has  $\sigma_{P_j} = \sum_{p \geq 0} b_p B_{[p]}$ , for some  $b_p \in \mathbb{C}$  and  $B \in V$ ; such vector  $B$  is obtained from  $A$  by the action

of an element in  $\text{Aut } \mathcal{O}$  producing the isomorphism between the stalks of  $\mathcal{V}_C$  at  $P_i$  and at  $P_j$ , hence  $B \in V_{\leq k}$ . It follows that  $\sigma_{P_j} \cdot M_{d_j}^j \subset M_{d_j+k-1}^j$ . From the identity

$$\sigma(m_1 \otimes \cdots \otimes m_n) = \sum_{j=1}^n m_1 \otimes \cdots \otimes \sigma_{P_j}(m_j) \otimes \cdots \otimes m_n,$$

one has

$$m_1 \otimes \cdots \otimes A_{(-l)}m_i \otimes \cdots \otimes m_n \in \mathcal{F}_{\sum_j d_j+k+l-2} M^\bullet + \mathcal{L}_{C \setminus P_\bullet}(V, D) \cdot M^\bullet.$$

Since the element on the left-hand side is in  $\mathcal{F}_{\sum_j d_j+k+l-1} M^\bullet$ , it follows that it maps to zero via  $\pi$ . The claim follows.

*Step 3.* After Step 2, the map  $\pi$  factors through

$$(26) \quad M^1/C_N(U, M^1) \otimes \cdots \otimes M^n/C_N(U, M^n) \xrightarrow{\pi} \text{gr}_\bullet \left( M_{\mathcal{L}_{C \setminus P_\bullet}(V, D)}^\bullet \right).$$

By [AN, Prop. 4.5], there is a positive integer  $k$  such that  $C_k(M^i) \subset C_N(U, M^i)$  for all  $i$ . In particular,  $\dim M^i/C_N(U, M^i) < \dim M^i/C_k(M^i)$ . These are finite as the  $M^i$  are all  $C_k$ -cofinite by [Buh]. It follows that the source in (26) is finite-dimensional, hence so is the target. This implies that the coinvariants are finite-dimensional as well.  $\square$

## 6. THE MODULES $Z$ AND $\bar{Z}$

We introduce here two modules which will aide the proof of the factorization property of spaces of coinvariants. In §6.1 we define the modules  $Z$  and  $\bar{Z}$  and give their properties, and in §6.2 we study spaces of coinvariants with a module  $\bar{Z}$ .

**6.1. Definitions and properties.** Let  $V$  be a conformal vertex algebra. Recall the associative algebra  $\mathcal{U}(V)$  from §1.4. Consider the  $\mathcal{U}(V)^{\otimes 2}$ -module

$$Z := \left( \text{Ind}_{\mathcal{U}(V)_{\leq 0}}^{\mathcal{U}(V)} A(V) \right)^{\otimes 2} = \left( \mathcal{U}(V) \otimes_{\mathcal{U}(V)_{\leq 0}} A(V) \right)^{\otimes 2}$$

where  $\mathcal{U}(V)_{<0}$  acts trivially on  $A(V)$ , and the action of  $\mathcal{U}(V)_0$  on  $A(V)$  is induced from the projection  $\mathcal{U}(V)_0 \rightarrow A(V)$ . With the notation from §1.6, one has that  $Z = M(A(V))^{\otimes 2}$ , where  $M(A(V))$  is the generalized Verma  $\mathcal{U}(V)$ -module induced from the natural representation  $A(V)$  of  $A(V)$ .

In §7, we will also consider a quotient  $\bar{Z}$  of  $Z$ . Let  $\mathcal{P}$  be the subalgebra of  $\mathcal{U}(V)^{\otimes 2}$  generated by  $\mathcal{U}(V) \otimes_{\mathbb{C}} \mathcal{U}(V)_{\leq 0}$  and  $\mathcal{U}(V)_{\leq 0} \otimes_{\mathbb{C}} \mathcal{U}(V)$ . Consider the  $\mathcal{U}(V)^{\otimes 2}$ -module

$$\bar{Z} := \text{Ind}_{\mathcal{P}}^{\mathcal{U}(V)^{\otimes 2}} A(V) = \left( \mathcal{U}(V)^{\otimes 2} \right) \otimes_{\mathcal{P}} A(V)$$

where  $\mathcal{U}(V) \otimes_{\mathbb{C}} \mathcal{U}(V)_{<0}$  and  $\mathcal{U}(V)_{<0} \otimes_{\mathbb{C}} \mathcal{U}(V)$  act trivially on  $A(V)$ , and the action of  $\mathcal{U}(V)_0 \otimes_{\mathbb{C}} \mathcal{U}(V)_0$  on  $A(V)$  is induced via the natural surjection  $\mathcal{U}(V)_0 \otimes_{\mathbb{C}} \mathcal{U}(V)_0 \rightarrow A(V) \otimes_{\mathbb{C}} A(V)$  from the action of  $A(V) \otimes_{\mathbb{C}} A(V)$  given by

$$(a \otimes b)(c) = a \cdot c \cdot (-\vartheta(b)), \quad \text{for } a \otimes b \in A(V) \otimes A(V), c \in A(V).$$

**Lemma 6.1.1.** *Let  $V$  be a rational conformal vertex algebra. One has  $\mathcal{U}(V)^{\otimes 2}$ -module isomorphisms*

$$Z \cong \bigoplus_{W, Y \in \mathcal{W}} (W \otimes W_0^\vee) \otimes (Y \otimes Y_0^\vee) \quad \text{and} \quad \bar{Z} \cong \bigoplus_{W \in \mathcal{W}} W \otimes W',$$

where  $\mathcal{W}$  is the set of all the finitely many simple  $V$ -modules.

*Proof.* Since  $V$  is rational, the algebra  $A(V)$  is semisimple [Zhu2]. From Wedderburn's theorem, one has  $A(V) = \bigoplus_{E \in \mathcal{E}} E \otimes E^\vee$ , where  $\mathcal{E}$  is the set of all finitely many simple  $A(V)$ -modules. Using the one-to-one correspondence between simple  $V$ -modules and simple  $A(V)$ -modules [Zhu2], and rationality of  $V$  which implies that the  $V$ -module induced from any simple  $A(V)$ -module is simple, it follows that each simple  $V$ -module is  $W = \mathcal{U}(V) \otimes_{\mathcal{U}(V)_{\leq 0}} E$ , for some  $E \in \mathcal{E}$ . Moreover, there exists a canonical  $V$ -module isomorphism  $\mathcal{U}(V) \otimes_{\mathcal{U}(V)_{\leq 0}} E^\vee \cong (\mathcal{U}(V) \otimes_{\mathcal{U}(V)_{\leq 0}} E)'$ , for  $E \in \mathcal{E}$  [NT, Prop. 7.2.1]. The statement follows by linearity.  $\square$

**6.2. Replacing coinvariants with the module  $Z$ .** In this section we describe a Lie subalgebra  $\mathcal{L}_{C \setminus P_\bullet}(V, \{Q_1, Q_2\})$  of  $\mathcal{L}_{C \setminus P_\bullet}(V)$ , which provides the following statement, generalizing [NT, Prop. 7.2.2, Cor. 8.6.2] from rational curves to curves of arbitrary genus.

**Proposition 6.2.1.** *Consider a coordinatized  $(n+2)$ -pointed smooth curve  $(C, P_\bullet \sqcup Q_\bullet, t_\bullet \sqcup s_\bullet)$ , possibly disconnected, such that  $C \setminus P_\bullet$  is affine. Let  $V$  be a rational conformal vertex algebra. Given  $V$ -modules  $M^1, \dots, M^n$ , the map*

$$M^\bullet \rightarrow M^\bullet \otimes Z, \quad w \mapsto w \otimes \mathbf{1}^{A(V)} \otimes \mathbf{1}^{A(V)},$$

where  $\mathbf{1}^{A(V)} \in A(V)$  is the unit, induces an isomorphism of vector spaces

$$h: M_{\mathcal{L}_{C \setminus P_\bullet}(V, \{Q_1, Q_2\})}^\bullet \xrightarrow{\cong} (M^\bullet \otimes_{\mathbb{C}} Z)_{\mathcal{L}_{C \setminus P_\bullet \sqcup Q_\bullet}(V)}.$$

The proof of Proposition 6.2.1, given after the definition of the Lie algebra  $\mathcal{L}_{C \setminus P_\bullet}(V, \{Q_1, Q_2\})$ , involves the study of the chiral Lie algebra in §3, Proposition 3.4.1, and an extension of the statement on propagation of vacua given in Remark 6.2.2.

As in the statement, let  $C$  be a smooth curve, possibly disconnected, with two nonempty, disjoint sets of distinct marked points  $P_\bullet = (P_1, \dots, P_n)$  and  $Q_\bullet = (Q_1, Q_2)$ . Assume that  $C \setminus P_\bullet$  is affine. After Lemmas 2.7.2 and 2.7.1, one has

$$H^0(C \setminus P_\bullet, \mathcal{V}_C) \cong \bigoplus_{k \geq 0} H^0(C \setminus P_\bullet, V_k \otimes_{\mathbb{C}} \omega_C^{\otimes -k}).$$

Using this isomorphism, consider the following Lie subalgebra of the chiral Lie algebra  $\mathcal{L}_{C \setminus P_\bullet}(V)$ :

(27)

$$\mathcal{L}_{C \setminus P_\bullet}(V, \{Q_1, Q_2\}) := \frac{\bigoplus_{k \geq 0} H^0(C \setminus P_\bullet, V_k \otimes_{\mathbb{C}} \omega_C^{\otimes 1-k}(-kQ_1 - kQ_2))}{\nabla H^0(C \setminus P_\bullet, \mathcal{V}_C)}.$$

As in (24),  $\nabla H^0(C \setminus P_\bullet, \mathcal{V}_C)$  is the intersection of  $\text{Im} \nabla$  with the subspace  $\bigoplus_{k \geq 0} H^0(C \setminus P_\bullet, V_k \otimes_{\mathbb{C}} \omega_C^{\otimes 1-k}(-kQ_1 - kQ_2))$  of  $H^0(C \setminus P_\bullet, \mathcal{V}_C \otimes \omega_C)$ .

Set  $\mathfrak{L}_{P_\bullet}(V) := \bigoplus_{i=1}^n \mathfrak{L}_{P_i}(V)$  and  $\mathfrak{L}_{Q_\bullet}(V) := \mathfrak{L}_{Q_1}(V) \oplus \mathfrak{L}_{Q_2}(V)$ . Fixing formal coordinates  $t_i$  at the point  $P_i$  and  $s_i$  at  $Q_i$ , one has Lie algebra homomorphisms

$$(28) \quad \mathcal{L}_{C \setminus P_\bullet}(V, \{Q_1, Q_2\}) \rightarrow \mathfrak{L}_{P_\bullet}(V) \text{ and } \mathcal{L}_{C \setminus P_\bullet}(V, \{Q_1, Q_2\}) \rightarrow \mathfrak{L}_{Q_\bullet}(V).$$

The Lie algebra  $\mathcal{L}_{C \setminus P_\bullet}(V, \{Q_1, Q_2\})$  consists of the elements in  $\mathcal{L}_{C \setminus P_\bullet}(V)$  whose image in  $\mathfrak{L}_{Q_\bullet}(V) \cong \mathfrak{L}(V)^{\oplus 2}$  via the restriction map in (28) lies in  $\mathfrak{L}(V)_{<0}^{\oplus 2} \subset \mathfrak{L}_{Q_\bullet}(V)$ . Indeed, the image of an element of  $\mathcal{L}_{C \setminus P_\bullet}(V, \{Q_1, Q_2\})$  in  $\mathfrak{L}_{Q_\bullet}(V)$  is a linear combination of elements of type

$$\sum_{i \geq k} a_i A_{[i]} \oplus \sum_{j \geq k} b_j B_{[j]} \in \mathfrak{L}(V)_{<0}^{\oplus 2}$$

for homogeneous  $A, B \in V$  of degree  $k \geq 0$  and coefficients  $a_i, b_j \in \mathbb{C}$ .

**Remark 6.2.2.** To prove Proposition 6.2.1 we are going to use a generalization of the statement on propagation of vacua from [FBZ, Thm 10.3.1], originally stated for affine Lie algebras in [TUY, Prop. 2.2.3], to slightly larger spaces of coinvariants and conformal blocks. Namely, one has a canonical isomorphism of vector spaces

$$\text{Hom}\left((M^\bullet \otimes V^\bullet)_{\mathcal{L}_{C \setminus P_\bullet \sqcup R_\bullet}(V, \{Q_1, Q_2\})}, \mathbb{C}\right) \cong \text{Hom}\left(M_{\mathcal{L}_{C \setminus P_\bullet}(V, \{Q_1, Q_2\})}^\bullet, \mathbb{C}\right)$$

for a set of distinct points  $R_\bullet$  disjoint from  $P_\bullet \sqcup Q_\bullet$ .

For this, let  $R \in C \setminus P_\bullet \sqcup Q_\bullet$  and fix a formal coordinate at  $R$ . As in [FBZ, §10.3], the natural map

$$\xi: \text{Hom}\left((M^\bullet \otimes V)_{\mathcal{L}_{C \setminus P_\bullet \sqcup R}(V, \{Q_1, Q_2\})}, \mathbb{C}\right) \rightarrow \text{Hom}\left(M_{\mathcal{L}_{C \setminus P_\bullet}(V, \{Q_1, Q_2\})}^\bullet, \mathbb{C}\right)$$

given by  $\xi(\varphi)(w) = \varphi(w \otimes \mathbf{1}^V)$  is an isomorphism. Indeed, using the strong residue theorem [FBZ, Thm 9.2.9], [Tat], we can identify the target of  $\xi$  as those linear functionals  $\varphi$  on  $M^\bullet$  such that for any  $w \in M^\bullet$ , the elements  $\langle \varphi, \mathcal{Y}^{M^i}(-)(w) \rangle$  of  $\bigoplus_{k \geq 0} H^0(D_{P_i}^\times, V_k^\vee \otimes \omega_C^{\otimes k}(kQ_1 + kQ_2))$ , with  $i = 1, \dots, n$ , can be extended to the same section  $\varphi_w$  of

$$\bigoplus_{k \geq 0} H^0(C \setminus P_\bullet, V_k^\vee \otimes \omega_C^{\otimes k}(kQ_1 + kQ_2)).$$

Here  $\mathcal{Y}^{M^i}(-)$  denotes the coordinate independent version of  $Y^{M^i}(-, z)$  as in [FBZ, §7.3]. Using this section, one realizes the inverse of  $\xi$  by

$$\xi^{-1}(\varphi)(w \otimes A) = \varphi_w(A).$$

One checks that this realizes the inverse of  $\xi$  as in the proof of [FBZ, §10.3.1]. Iterating on the number of points in  $R_\bullet$  gives the assertion.

*Proof of Proposition 6.2.1.*

*Step 1.* We first show that the map  $h$  is well-defined. Observe that  $Z$  is naturally equipped with a left action of  $\mathcal{L}_{C \setminus P_\bullet \sqcup Q_\bullet}(V)$  induced by the Lie

algebra homomorphisms  $\mathcal{L}_{C \setminus P_\bullet \sqcup Q_\bullet}(V) \rightarrow \mathfrak{L}_{Q_\bullet}(V) \rightarrow \mathcal{U}(V)^{\otimes 2}$ . For  $\sigma \in \mathcal{L}_{C \setminus P_\bullet}(V, \{Q_1, Q_2\}) \subset \mathcal{L}_{C \setminus P_\bullet \sqcup Q_\bullet}(V)$ , let  $\sigma_{P_\bullet}$  be the image of  $\sigma$  in  $\mathfrak{L}_{P_\bullet}(V)$  via the restriction map in (28), and similarly let  $\sigma_{Q_i}$  be its image in  $\mathfrak{L}_{Q_i}(V)$ , for  $i = 1, 2$ . Since  $\sigma_{Q_i} \in \mathfrak{L}(V)_{<0}$ , the elements  $\sigma_{Q_1} \otimes 1$  and  $1 \otimes \sigma_{Q_2}$  act trivially on  $A(V) \otimes A(V) \subset Z$ . This implies

$$(29) \quad \begin{aligned} \sigma_{P_\bullet}(w) \otimes \mathbf{1}^{A(V)} \otimes \mathbf{1}^{A(V)} &= \\ \sigma(w \otimes \mathbf{1}^{A(V)} \otimes \mathbf{1}^{A(V)}) - w \otimes \sigma_{Q_1}(\mathbf{1}^{A(V)}) \otimes \mathbf{1}^{A(V)} - w \otimes \mathbf{1}^{A(V)} \otimes \sigma_{Q_2}(\mathbf{1}^{A(V)}) & \\ &= \sigma(w \otimes \mathbf{1}^{A(V)} \otimes \mathbf{1}^{A(V)}) \end{aligned}$$

for  $w \in M^\bullet$ . It follows that the zero element is mapped to the zero element, hence the map  $h$  between the spaces of coinvariants is well-defined.

*Step 2.* Next, we show that the map  $h$  is surjective: given  $w \otimes z_1 \otimes z_2$  in  $M^\bullet \otimes Z$ , there exists  $w' \in M^\bullet$  such that

$$w \otimes z_1 \otimes z_2 \equiv w' \otimes \mathbf{1}^{A(V)} \otimes \mathbf{1}^{A(V)} \pmod{\mathcal{L}_{C \setminus P_\bullet \sqcup Q_\bullet}(V)(M^\bullet \otimes Z)}.$$

By linearity, and by reordering elements in  $\mathcal{U}(V)$ , we can reduce to the case

$$z_1 \otimes z_2 = D_l \cdots D_1 \mathbf{1}^{A(V)} \otimes E_m \cdots E_1 \mathbf{1}^{A(V)},$$

with each  $D_i$  and  $E_j$  in  $\mathfrak{L}(V)_{\geq 0}$ . The surjectivity is clear when  $l = m = 0$ . By induction on  $l$  (and similarly on  $m$ ), it is then enough to show that

$$w \otimes z_1 \otimes z_2 \equiv w' \otimes z'_1 \otimes z_2 \pmod{\mathcal{L}_{C \setminus P_\bullet \sqcup Q_\bullet}(V)(M^\bullet \otimes Z)}$$

for some  $w'$  in  $M^\bullet$ , when  $z_1 = D_{[d]}(z'_1)$  for some homogeneous  $D \in V$  and  $D_{[d]}$  in  $\mathfrak{L}(V)_{\geq 0}$ . Each component of the curve  $C$  has at least one of the marked points in  $P_\bullet$ . By Proposition 3.4.1, there exists  $\sigma \in \mathcal{L}_{C \setminus P_\bullet \sqcup Q_\bullet}(V)$  such that

$$\begin{aligned} \sigma_{Q_1} \equiv D_{[d]} &\in \mathfrak{L}_{Q_1}(V)/\mathfrak{L}_{Q_1}(V, NQ_1), \\ \sigma_{Q_2} \equiv 0 &\in \mathfrak{L}_{Q_2}(V)/\mathfrak{L}_{Q_2}(V, NQ_2), \end{aligned}$$

for  $N \gg 0$ . It is enough to take  $N$  such that both

$$D_{[N]}(z'_1) \otimes z_2 \quad \text{and} \quad z'_1 \otimes D_N(z_2)$$

are zero in  $Z$ . Such  $N$  exists because  $\mathcal{U}(V)$  acts smoothly on each component of  $Z$ . Moreover,  $\sigma_{Q_2}$  is  $\sum_{i \geq N} a_i A_{[i]}$ , with  $a_i \in \mathbb{C}$  and  $A \in V$  obtained from  $D$  via the action of an element of  $\text{Aut } \mathcal{O}$  producing the isomorphism between the stalks of  $\mathcal{V}_C$  at  $Q_1$  and at  $Q_2$ . It follows that  $A \in V_{\leq \deg D}$ , hence  $\sigma_{Q_2}(z_2) = 0$ . This implies

$$\sigma_{Q_1}(z'_1) \otimes z_2 + z'_1 \otimes \sigma_{Q_2}(z_2) = D_{[d]} \cdot z'_1 \otimes z_2 = z_1 \otimes z_2.$$

It follows that

$$w \otimes z_1 \otimes z_2 = \sigma(w \otimes z'_1 \otimes z_2) - \sigma_{P_\bullet}(w) \otimes z'_1 \otimes z_2,$$

hence

$$w \otimes z_1 \otimes z_2 \equiv -\sigma_{P_\bullet}(w) \otimes z'_1 \otimes z_2, \pmod{\mathcal{L}_{C \setminus P_\bullet \sqcup Q_\bullet}(V)(M^\bullet \otimes Z)}.$$

Repeating the same argument for  $z_2$ , the surjectivity of  $h$  follows.

*Step 3.* Finally, we show that  $h$  is injective. Equivalently, we show that the dual map

$$h^\vee : \text{Hom} \left( (M^\bullet \otimes_{\mathbb{C}} Z)_{\mathcal{L}_{C \setminus P_\bullet \sqcup Q_\bullet}(V)}, \mathbb{C} \right) \rightarrow \text{Hom} \left( M^\bullet_{\mathcal{L}_{C \setminus P_\bullet}(V, \{Q_1, Q_2\})}, \mathbb{C} \right)$$

given by

$$h^\vee(\Phi)(w) = \Phi \left( w \otimes \mathbf{1}^{A(V)} \otimes \mathbf{1}^{A(V)} \right), \quad \text{for } w \in M^\bullet,$$

is surjective. Let  $\Phi_{0,0}$  be an element in the target of  $h^\vee$ , i.e.,  $\Phi_{0,0}$  is a linear functional on  $M^\bullet$  vanishing on the subspace  $\mathcal{L}_{C \setminus P_\bullet}(V, \{Q_1, Q_2\}) \cdot M^\bullet$ . In the following, we construct a linear functional  $\Phi$  on  $M^\bullet \otimes_{\mathbb{C}} Z$  vanishing on  $\mathcal{L}_{C \setminus P_\bullet \sqcup Q_\bullet}(V) \cdot (M^\bullet \otimes_{\mathbb{C}} Z)$  such that  $h^\vee(\Phi) = \Phi_{0,0}$ . The argument uses the formalism of systems of correlation functions, as developed in this setting in [NT, §5] for rational curves, with the extensions to curves of arbitrary genus in [FBZ, §10.3] (see Remark 4.2.1).

*Step 3(a).* Given  $\Phi_{0,0}$ , one first constructs linear functionals  $\Phi_{l,m}$  on  $M^\bullet \otimes V^{\otimes l+m}$  for all  $l, m \geq 0$ . By induction on  $l$  and  $m$ , it is enough to construct  $\Phi_{l,m}$  starting from  $\Phi_{l-1,m}$ . This uses a version of propagation of vacua similar to the one proved in [FBZ, §10.3.1]. Given  $l, m \in \mathbb{Z}_{\geq 0}$ , fix points  $R_\bullet = R_{[l]} = (R_1, \dots, R_l)$  and  $S_\bullet = (S_1, \dots, S_m)$  in  $C \setminus P_\bullet \sqcup Q_\bullet$  such that  $R_\bullet \sqcup S_\bullet$  is a collection of distinct points. Additionally, fix formal coordinates at the points  $R_\bullet \sqcup S_\bullet$ . By the generalization of propagation of vacua given in Remark 6.2.2, there is a canonical isomorphism of vector spaces

$$\begin{aligned} \xi : \text{Hom} \left( (M^\bullet \otimes_{\mathbb{C}} V^{\otimes l+m})_{\mathcal{L}_{C \setminus P_\bullet \sqcup R_{[l]} \sqcup S_\bullet}(V, \{Q_1, Q_2\})}, \mathbb{C} \right) \\ \rightarrow \text{Hom} \left( (M^\bullet \otimes_{\mathbb{C}} V^{\otimes l-1+m})_{\mathcal{L}_{C \setminus P_\bullet \sqcup R_{[l-1]} \sqcup S_\bullet}(V, \{Q_1, Q_2\})}, \mathbb{C} \right). \end{aligned}$$

One then defines

$$(30) \quad \Phi_{l,m} := \xi^{-1}(\Phi_{l-1,m}).$$

Observe that we could have equivalently defined  $\Phi_{l,m} = \xi^{-1}(\Phi_{l,m-1})$  interchanging the role of  $l$  and  $m$ .

*Step 3(b).* The linear functional  $\Phi_{l,m}$  gives rise to a meromorphic form on  $C^{l+m}$  as follows. For fixed  $l$  and  $m$ , by varying the points  $R_\bullet$  and  $S_\bullet$  in  $C \setminus P_\bullet \sqcup Q_\bullet$ , the space of conformal blocks, as in the source of the map  $\xi$  above, becomes a fiber of the sheaf of conformal blocks  $\mathbb{V}(M^\bullet \otimes V^{l+m})^\vee$  on  $C^{l+m} \setminus \Delta \cup \Gamma_{P_\bullet} \cup \Gamma_{Q_\bullet}$ . Here  $\Delta$  is the locus of  $(l+m)$ -tuples where two components coincide;  $\Gamma_{P_\bullet}$  is the locus of  $(l+m)$ -tuples where one component equals one of the points in  $P_\bullet$ ; and  $\Gamma_{Q_\bullet}$  is the locus of  $(l+m)$ -tuples where one component equals  $Q_1$  or  $Q_2$ . Similar sheaves have been studied in [FBZ, DGT2]. The

linear functionals  $\Phi_{l,m}$  patch together to define a global meromorphic section  $\Phi_{l,m(R_\bullet, S_\bullet)}$  of  $\mathbb{V}(M^\bullet \otimes V^{l+m})^\vee$ .

Now fix an element  $w \otimes D^\bullet \otimes E^\bullet \in M \otimes V^{\otimes l+m}$ , with  $D^\bullet = D^l \otimes \cdots \otimes D^1$  and  $E^\bullet = E^m \otimes \cdots \otimes E^1$ . As in Remark 6.2.2, the strong residue theorem allows us to attach to such a vector and to the section  $\Phi_{l,m(R_\bullet, S_\bullet)}$  a meromorphic form on  $C^{l+m}$  denoted by

$$(31) \quad \Phi_{l,m}(w \otimes D^\bullet \otimes E^\bullet)_{(R_\bullet, S_\bullet)}.$$

When the elements  $D^i$  and  $E^j$  are homogeneous of degree  $d_i$  and  $e_j$ , respectively, this form is an element of

$$H^0 \left( C^{l+m} \setminus \Delta \cup \Gamma_{P_\bullet}, \omega_C^{\boxtimes_{i=1}^l d_i} \boxtimes \omega_C^{\boxtimes_{j=1}^m e_j} \left( \sum_{i=1}^l d_i \Gamma_{R_i, Q_\bullet} + \sum_{j=1}^m e_j \Gamma_{S_j, Q_\bullet} \right) \right),$$

where  $\Gamma_{R_i, Q_\bullet} \subset C^{l+m}$  is the locus where  $R_i$  coincides with  $Q_1$  or  $Q_2$ , and  $\Gamma_{S_j, Q_\bullet} \subset C^{l+m}$  is the locus where  $S_j$  coincides with  $Q_1$  or  $Q_2$ . Note that the order of the poles along  $\Delta$  and  $\Gamma_{P_\bullet}$  is unbounded.

*Step 3(c).* In the following, it is enough to consider the restriction of (31) to a rational subset of  $C^{l+m}$ . Namely, consider a rational subset  $U \subset C \setminus Q_\bullet$  containing the points  $P_\bullet$ . After removing finitely many points in  $C \setminus Q_\bullet$ , one can indeed find such a set: consider a ramified cover  $C \rightarrow \mathbb{P}^1$  and remove enough points to get an étale map  $U \rightarrow \mathbb{A}^1$ . Such a map induces a formal coordinate at every point of  $U$ , hence a trivialization of  $\mathcal{A}ut_C$  on  $U$  [FBZ, §6.5.2]. That is, there exists a global coordinate  $t$  on  $U$  inducing a formal coordinate  $t - u$  at every point  $u \in U$ . In turn, this gives a trivialization of  $\mathcal{V}_C$  on  $U$ . In particular, using coordinates  $t - x_i$  at  $R_i$  and  $t - y_j$  at  $S_j$ , and given homogeneous  $E^i$  and  $D^j$  in  $V$  of degree  $d_i$  and  $e_j$ , respectively, we can rewrite the restriction of (31) to  $U^{l+m}$  as

$$(32) \quad \bar{\Phi}_{l,m}(w \otimes D^\bullet \otimes E^\bullet)_{(R_\bullet, S_\bullet)} (dx_1)^{d_1} \cdots (dx_l)^{d_l} (dy_1)^{e_1} \cdots (dy_m)^{e_m}$$

where  $\bar{\Phi}_{l,m}(w \otimes D^\bullet \otimes E^\bullet)_{(R_\bullet, S_\bullet)}$  is a rational function with poles along  $\Delta$  and  $\Gamma_{P_\bullet}$ .

Note that we require the points  $Q_1$  and  $Q_2$  to lie outside of  $U$ . The formal coordinate  $s_i$  at  $Q_i$  satisfies  $s_i = \rho_i(t)$ , for a change of variable  $\rho_i \in \text{Aut}(\mathcal{O})$ , for  $i = 1, 2$ . This element  $\rho_i$  induces an isomorphism  $\mathfrak{L}_t(V) \rightarrow \mathfrak{L}_{s_i}(V)$ , still denoted  $\rho_i$ .

*Step 3(d).* We show that the set of linear functionals  $\{\Phi_{l,m}\}_{l,m \geq 0}$  gives rise to a linear functional  $\Phi$  on  $M^\bullet \otimes Z$ . Recall that the module  $Z$  is linearly generated by

$$z_1 \otimes z_2 = \rho_1(D_{[i_1]}^l) \cdots \rho_1(D_{[i_1]}^1) \mathbf{1}^{A(V)} \otimes \rho_2(E_{[j_m]}^m) \cdots \rho_2(E_{[j_1]}^1) \mathbf{1}^{A(V)}$$

with  $D^1, \dots, D^l, E^1, \dots, E^m \in V$ , and  $i_1, \dots, i_l, j_1, \dots, j_m \in \mathbb{Z}$ . For such  $z_1 \otimes z_2$  and  $w \in M^\bullet$ , the linear functional  $\Phi$  is defined as

$$\Phi(w \otimes z_1 \otimes z_2) := \left( \frac{1}{2\pi\sqrt{-1}} \right)^{l+m} \oint_{|y_m|=r_{l+m}} \cdots \oint_{|y_1|=r_{l+1}} \oint_{|x_l|=r_l} \cdots \oint_{|x_1|=r_1} \bar{\Phi}_{l,m}(w \otimes D^\bullet \otimes E^\bullet)_{(R_\bullet, S_\bullet)} x_1^{i_1} \cdots x_l^{i_l} y_1^{j_1} \cdots y_m^{j_m} dx_1 \cdots dx_l dy_1 \cdots dy_m$$

where  $r_1 > \cdots > r_{l+m} > \max_i |P_i|$  on the rational subset  $U$ . This implies that all the  $P_i$ 's are contained inside the region bounded by the interval of integration, and that the interval of integration avoids the locus  $\Delta \cup \Gamma_{P_\bullet} \subset U^{l+m}$ . The above defines  $\Phi$  on the set of generators of  $M^\bullet \otimes Z$  and by construction

$$\Phi(w \otimes \mathbf{1}^{A(V)} \otimes \mathbf{1}^{A(V)}) = \Phi_{0,0}(w).$$

*Step 3(e).* We are left to prove that such  $\Phi$  is compatible with the relations defining  $Z$ , and vanishes on  $\mathcal{L}_{C \setminus P_\bullet \sqcup Q_\bullet}(V) \cdot (M^\bullet \otimes_{\mathbb{C}} Z)$ .

Recall that  $Z$  is spanned by  $z_1 \otimes z_2$  as above modulo some relations corresponding to the weak associativity of  $V$  as a  $V$ -module. Since the linear functionals  $\Phi_{l,m}$ , and hence  $\Phi$ , are defined in terms of meromorphic sections arising from the vertex operators  $Y^{M^i}(-, z)$ , it follows that the relations among elements of  $Z$  are preserved, hence  $\Phi$  is indeed a linear functional on  $M^\bullet \otimes Z$ .

Finally, we verify that  $\Phi$  vanishes on  $\mathcal{L}_{C \setminus P_\bullet \sqcup Q_\bullet}(V) (M^\bullet \otimes_{\mathbb{C}} Z)$ . For this, fix  $\sigma$  in  $\mathcal{L}_{C \setminus P_\bullet \sqcup Q_\bullet}(V)$ . On the affine  $C \setminus P_\bullet \sqcup Q_\bullet$ , the section  $\sigma$  can be described as the tensor product of a section  $\mathcal{B}$  of  $\mathcal{V}_C$  and a 1-differential  $\mu$ . One can choose a rational subset  $U \subset C \setminus Q_\bullet$  as in *Step 3(c)* such that  $\mathcal{V}_C$  can be trivialized on  $U$ . Therefore, the restriction  $\sigma_{P_i}$  of  $\sigma$  from  $C \setminus P_\bullet \sqcup Q_\bullet$  to the punctured formal disk  $D_{P_i}^\times$  can be described as

$$A \otimes \mu_{P_i}, \quad \text{for some } A \in V \text{ independent of } P_i,$$

where  $\mu_{P_i}$  is the Laurent series expansion of  $\mu$  at  $P_i$ . Similarly, after trivializing  $\mathcal{V}_C$  on a rational subset containing the points  $Q_\bullet$ , the restriction  $\sigma_{Q_i}$  of  $\sigma$  to  $D_{Q_i}^\times$  can be described as  $B \otimes \mu_{Q_i}$ , for some  $B \in V$  independent of  $Q_i$ . For  $w \otimes z_1 \otimes z_2 \in M^\bullet \otimes Z$ , one has

$$\begin{aligned} (33) \quad & \Phi(w \otimes \sigma_{Q_1}(z_1) \otimes z_2) + \Phi(w \otimes z_1 \otimes \sigma_{Q_2}(z_2)) \\ &= \Phi(w \otimes \text{Res}_{s_1=0} Y(B, s_1) z_1 \mu_{Q_1} \otimes z_2) + \Phi(w \otimes z_1 \otimes \text{Res}_{s_2=0} Y(B, s_2) z_2 \mu_{Q_2}) \\ &= \text{Res}_{s_1=0} \Phi(w \otimes Y(B, s_1) z_1 \otimes z_2) \mu_{Q_1} + \text{Res}_{s_2=0} \Phi(w \otimes z_1 \otimes Y(B, s_2) z_2) \mu_{Q_2}. \end{aligned}$$

From the definition of  $\Phi$  in terms of the linear functionals  $\{\Phi_{l,m}\}_{l,m}$ , and the identity  $\Phi_{l+1,m} = \Phi_{l,m+1}$  one has that

$$\Phi(w \otimes Y(B, s)(z_1) \otimes z_2) = \Phi(w \otimes z_1 \otimes Y(B, s)(z_2))$$

for some formal variable  $s$ . In particular,

$$\Phi(w \otimes Y(B, s_1)(z_1) \otimes z_2) \mu_{Q_1} \quad \text{and} \quad \Phi(w \otimes z_1 \otimes Y(B, s_2)(z_2)) \mu_{Q_2}$$

are the Laurent series expansions at  $Q_1$  and  $Q_2$ , respectively, of the meromorphic 1-differential on  $C$ , with poles at the points  $P_\bullet \sqcup Q_\bullet$  and regular elsewhere, given by

$$\Phi(w \otimes \mathcal{Y}(\mathcal{B})(z_1) \otimes z_2) \mu,$$

where  $\Phi(w \otimes \mathcal{Y}(\mathcal{B})(z_1) \otimes z_2)$  is the regular function on  $C \setminus P_\bullet \sqcup Q_\bullet$  whose fiber over a point  $T$  with formal coordinate  $t$  is given by  $\Phi(w \otimes Y(\mathcal{B}|_T, t)(z_1) \otimes z_2)$ . From the residue theorem and the fact that  $\mathcal{B}|_{P_i} = A$ , it follows that (33) is equal to

$$-\sum_{i=1}^n \operatorname{Res}_{t_i=0} \Phi(w \otimes Y(A, t_i)(z_1) \otimes z_2) \mu_{P_i}.$$

By definition

$$\Phi(\sigma_{P_\bullet}(w) \otimes z_1 \otimes z_2) = \sum_{i=1}^n \operatorname{Res}_{t_i=0} \Phi(Y(A, t_i)(w) \otimes z_1 \otimes z_2) \mu_{P_i},$$

where

$$Y(A, t_i)(w) := w_1 \otimes \cdots \otimes Y(A, t_i)(w_i) \otimes \cdots \otimes w_n$$

for  $w = w_1 \otimes \cdots \otimes w_n$ . Thus we are left to show that

$$\operatorname{Res}_{t_i=0} \Phi(w \otimes Y(A, t_i)(z_1) \otimes z_2) \mu_{P_i} = \operatorname{Res}_{t_i=0} \Phi(Y(A, t_i)(w) \otimes z_1 \otimes z_2) \mu_{P_i}$$

for each  $i$ . To verify this, fix  $w \otimes A \otimes v \in M^\bullet \otimes V^{\otimes l+m+1}$ . The left-hand side is computed by means of  $\Phi_{l+1,m}(w \otimes A \otimes v)$ , while the right-hand side by means of  $\Phi_{l,m}(Y(A, t_i)(w) \otimes v)$ . Considering the point corresponding to the element  $A$  to be in  $D_{P_i}^\times$  and fixing the points corresponding to the element  $v \in V^{\otimes l+m}$ ,  $\Phi_{l+1,m}(w \otimes A \otimes v)$  becomes a meromorphic function on  $D_{P_i}$ , denoted  $\Phi_{l+1,m}(w \otimes A \otimes v)_{(t_i)}$ . The desired identity follows from

$$\Phi_{l+1,m}(w \otimes A \otimes v)_{(t_i)} = \Phi_{l,m}(Y(A, t_i)(w) \otimes v)$$

which holds from propagation of vacua (this is similar to [FBZ, §10.3.2]).

The above argument gives

$$\Phi(w \otimes \sigma_{Q_1}(z_1) \otimes z_2) + \Phi(w \otimes z_1 \otimes \sigma_{Q_2}(z_2)) = -\Phi(\sigma_{P_\bullet}(w) \otimes z_1 \otimes z_2).$$

It follows that  $\Phi$  vanishes on  $\sigma(M^\bullet \otimes_{\mathbb{C}} Z)$  for all  $\sigma \in \mathcal{L}_{C \setminus P_\bullet \sqcup Q_\bullet}(V)$ , hence  $\Phi$  lies in the source of  $h^\vee$ .  $\square$

## 7. PROOF OF THEOREM 1

Here we prove our main result, which we state in complete detail below. For this let us first set some notation. Let  $(C, P_\bullet)$  be a stable  $n$ -pointed curve with exactly one node, denoted  $Q$ . Let  $\tilde{C} \rightarrow C$  be the normalization of  $C$ , let  $Q_+$  and  $Q_- \in \tilde{C}$  be the two preimages of  $Q$ , and set  $Q_\bullet = (Q_+, Q_-)$ . The curve  $\tilde{C}$  may not be connected. Suppose  $M^1, \dots, M^n$  are  $V$ -modules,

set  $M^\bullet = \otimes_{i=1}^n M^i$ , and let  $\mathscr{W}$  be the set of all simple  $V$ -modules. Consider the map

$$(34) \quad M^\bullet \rightarrow \bigoplus_{W \in \mathscr{W}} M^\bullet \otimes_{\mathbb{C}} W \otimes_{\mathbb{C}} W', \quad u \mapsto \bigoplus_{W \in \mathscr{W}} u \otimes \mathbf{1}^{W_0}$$

where  $\mathbf{1}^{W_0} = \text{id}_{W_0} \in \text{End}(W_0) \cong W_0 \otimes W_0^\vee$ . Here  $W_0$  is the degree zero space of the module  $W = \bigoplus_{i \geq 0} W_i$ . Recall that, by definition, the vector spaces  $W_0$  and  $W_0^\vee$  are finite-dimensional.

**The Factorization Theorem.** *Let  $V = \bigoplus_{i \geq 0} V_i$  be a rational,  $C_2$ -cofinite conformal vertex algebra with  $V_0 \cong \mathbb{C}$ . The map (34) gives rise to a canonical isomorphism of vector spaces*

$$\mathbb{V}(V; M^\bullet)_{(C, P_\bullet)} \cong \bigoplus_{W \in \mathscr{W}} \mathbb{V}(V; M^\bullet \otimes_{\mathbb{C}} W \otimes_{\mathbb{C}} W')_{(\tilde{C}, P_\bullet \sqcup Q_\bullet)}.$$

The proof we give here roughly follows the outline of the proof in [NT, §8.6], with the generalizations to coinvariants defined using the chiral Lie algebra instead of Zhu's Lie algebra, and for curves of arbitrary genus, made possible by Propositions 3.3.1 and 6.2.1.

*Proof.* By definition (23), due to propagation of vacua, we can reduce to the case  $C \setminus P_\bullet$  affine, after possibly adding more marked points  $P_i$  and corresponding modules  $V$ . Fix formal coordinates  $t_i$  at  $P_i$ , for each  $i = 1, \dots, n$ , and  $s_\pm$  at  $Q_\pm$ , so that we have Lie algebra homomorphisms

$$\mathcal{L}_{C \setminus P_\bullet}(V) \rightarrow \mathfrak{L}_{P_\bullet}(V) \quad \text{and} \quad \mathcal{L}_{\tilde{C} \setminus P_\bullet \sqcup Q_\bullet}(V) \rightarrow \mathfrak{L}_{P_\bullet}(V) \oplus \mathfrak{L}_{Q_\bullet}(V).$$

In the following, we show that (34) induces a canonical isomorphism

$$(35) \quad M^\bullet_{\mathcal{L}_{C \setminus P_\bullet}(V)} \cong \bigoplus_{W \in \mathscr{W}} (M^\bullet \otimes_{\mathbb{C}} W \otimes_{\mathbb{C}} W')_{\mathcal{L}_{\tilde{C} \setminus P_\bullet \sqcup Q_\bullet}(V)}.$$

Being independent of the choice of formal coordinates  $t_i$  and  $s_\pm$ , this implies the assertion made in Theorem 1.

We will argue that there is a commutative diagram

$$\begin{array}{ccc} M^\bullet_{\mathcal{L}_{\tilde{C} \setminus P_\bullet}(V, \{Q_+, Q_-\})} & \xrightarrow[\cong]{h} & (M^\bullet \otimes_{\mathbb{C}} Z)_{\mathcal{L}_{\tilde{C} \setminus P_\bullet \sqcup Q_\bullet}(V)} \\ \downarrow & & \downarrow \\ M^\bullet_{\mathcal{L}_{C \setminus P_\bullet}(V)} & \xrightarrow[\cong]{f} & (M^\bullet \otimes_{\mathbb{C}} \bar{Z})_{\mathcal{L}_{C \setminus P_\bullet \sqcup Q_\bullet}(V)}. \end{array}$$

Then after Lemma 6.1.1, the isomorphism  $f$  gives (35).

*Step 1.* The top horizontal isomorphism  $h$  is given by Proposition 6.2.1.

*Step 2.* We argue that there is an inclusion

$$\iota: \mathcal{L}_{\tilde{C} \setminus P_\bullet}(V, \{Q_+, Q_-\}) \hookrightarrow \mathcal{L}_{C \setminus P_\bullet}(V).$$

Indeed, by Proposition 3.3.1 an element of  $\mathcal{L}_{C \setminus P_\bullet}(V)$  can be realized as  $\sigma$  in  $\mathcal{L}_{\tilde{C} \setminus P_\bullet \sqcup Q_\bullet}(V)$  such that  $\sigma_{Q_\pm} \in \mathfrak{L}(V)_{\leq 0}$  and the restrictions  $[\sigma_{Q_\pm}]_0$  of  $\sigma_{Q_\pm}$  to  $\mathfrak{L}(V)_0$  satisfy  $[\sigma_{Q_-}]_0 = \vartheta [\sigma_{Q_+}]_0$ . In particular,  $\mathcal{L}_{\tilde{C} \setminus P_\bullet}(V, \{Q_+, Q_-\})$  from (27) is a Lie subalgebra of  $\mathcal{L}_{C \setminus P_\bullet}(V)$ , since its elements satisfy  $\sigma_{Q_\pm} \in \mathfrak{L}(V)_{< 0}$ .

*Step 3.* To show that the bottom horizontal map  $f$  is an isomorphism, it remains to verify that the kernel of the two vertical maps coincide.

*Step 3(a).* The kernel of the left vertical map is the space

$$K = \mathcal{L}_{C \setminus P_\bullet}(V) \left( M_{\mathcal{L}_{\tilde{C} \setminus P_\bullet}(V, \{Q_+, Q_-\})}^\bullet \right).$$

Note that for  $\sigma$  in  $\mathcal{L}_{C \setminus P_\bullet}(V) / \mathcal{L}_{\tilde{C} \setminus P_\bullet}(V, \{Q_+, Q_-\})$ , the formula for the bracket (4) gives

$$\left[ \varphi(\sigma), \varphi \left( \mathcal{L}_{\tilde{C} \setminus P_\bullet}(V, \{Q_+, Q_-\}) \right) \right] \subset \varphi \left( \mathcal{L}_{\tilde{C} \setminus P_\bullet}(V, \{Q_+, Q_-\}) \right),$$

where  $\varphi$  is as in (14). It follows that  $\sigma$  acts on the source of  $h$ . The left vertical map is thus the quotient by the action of  $\mathcal{L}_{C \setminus P_\bullet}(V) / \mathcal{L}_{\tilde{C} \setminus P_\bullet}(V, \{Q_+, Q_-\})$ .

*Step 3(b).* We conclude the argument by showing that the right vertical map coincides with the quotient by  $h(K)$ . Recall that  $h(w) = w \otimes \mathbf{1}^{A(V)} \otimes \mathbf{1}^{A(V)}$ , for  $w \in M^\bullet$ . For  $\sigma$  in  $\mathcal{L}_{C \setminus P_\bullet}(V)$ , one has

$$\begin{aligned} \sigma_{P_\bullet}(w) \otimes \mathbf{1}^{A(V)} \otimes \mathbf{1}^{A(V)} &\equiv -w \otimes \sigma_{Q_+} \left( \mathbf{1}^{A(V)} \right) \otimes \mathbf{1}^{A(V)} \\ &\quad - w \otimes \mathbf{1}^{A(V)} \otimes \sigma_{Q_-} \left( \mathbf{1}^{A(V)} \right) \pmod{\mathcal{L}_{\tilde{C} \setminus P_\bullet \sqcup Q_\bullet}(V) (M^\bullet \otimes Z)}. \end{aligned}$$

Since the elements of  $\mathcal{U}(V)_{< 0}$  act trivially on  $A(V)$  this is congruent to

$$-w \otimes [\sigma_{Q_+}]_0 \left( \mathbf{1}^{A(V)} \right) \otimes \mathbf{1}^{A(V)} - w \otimes \mathbf{1}^{A(V)} \otimes \vartheta [\sigma_{Q_+}]_0 \left( \mathbf{1}^{A(V)} \right)$$

modulo  $\mathcal{L}_{\tilde{C} \setminus P_\bullet \sqcup Q_\bullet}(V) (M^\bullet \otimes Z)$ . From lemma 6.1.1, one has

$$\begin{aligned} (M^\bullet \otimes Z)_{\mathcal{L}_{\tilde{C} \setminus P_\bullet \sqcup Q_\bullet}(V)} &\cong \bigoplus_{W, Y \in \mathscr{W}} \frac{M^\bullet \otimes W \otimes W_0^\vee \otimes Y \otimes Y_0^\vee}{\mathcal{L}_{\tilde{C} \setminus P_\bullet \sqcup Q_\bullet}(V) (M^\bullet \otimes W \otimes W_0^\vee \otimes Y \otimes Y_0^\vee)} \\ &\cong \bigoplus_{W, Y \in \mathscr{W}} (M^\bullet \otimes W \otimes Y)_{\mathcal{L}_{\tilde{C} \setminus P_\bullet \sqcup Q_\bullet}(V)} \otimes W_0^\vee \otimes Y_0^\vee. \end{aligned}$$

Recall Lemma 1.8.1 on the action of  $\mathfrak{L}(V)$  on  $W_0^\vee$ . It follows that the image of  $K$  in  $(M^\bullet \otimes Z)_{\mathcal{L}_{\tilde{C} \setminus P_\bullet \sqcup Q_\bullet}(V)}$  is

$$\bigoplus_{W, Y \in \mathscr{W}} (M^\bullet \otimes W \otimes Y)_{\mathcal{L}_{\tilde{C} \setminus P_\bullet \sqcup Q_\bullet}(V)} \otimes \mathcal{I} (W_0^\vee, Y_0^\vee)$$

with  $\mathcal{I} (W_0^\vee, Y_0^\vee) \subset W_0^\vee \otimes Y_0^\vee$  linearly spanned by

$$\psi_W \circ \vartheta (A_{[k-1]}) \otimes \psi_Y + \psi_W \otimes \psi_Y \circ A_{[k-1]},$$

where  $\psi_W \in W_0^\vee$ ,  $\psi_Y \in Y_0^\vee$  for  $W, Y \in \mathscr{W}$ , and  $A \in V_k$ , for  $k \geq 0$ . One has

$$W_0^\vee \otimes Y_0^\vee / \mathcal{I} (W_0^\vee, Y_0^\vee) = \text{Hom}_{A(V)} (W_0, Y_0^\vee),$$

and by Schur's Lemma, this is isomorphic to  $\mathbb{C}$  when  $Y = W'$  and zero otherwise. This and the description of  $\bar{Z}$  from Lemma 6.1.1 imply that, after taking the quotient of  $(M^\bullet \otimes_{\mathbb{C}} Z)_{\mathcal{L}_{\mathcal{C} \setminus P_\bullet \sqcup Q_\bullet}(V)}$  by the kernel of the left vertical map, one obtains  $(M^\bullet \otimes_{\mathbb{C}} \bar{Z})_{\mathcal{L}_{\mathcal{C} \setminus P_\bullet \sqcup Q_\bullet}(V)}$ , hence the statement.  $\square$

## 8. SEWING AND LOCAL FREENESS

In this section we prove Theorem 2. For this, we start with Theorem 8.3.1, a refined version of the Factorization Theorem. This requires the notion of formal smoothings, reviewed below.

**8.1. Formal smoothings.** For a  $\mathbb{C}$ -algebra  $R$  with smooth  $\text{Spec}(R)$ , let  $\mathcal{C}_0 \rightarrow S_0 = \text{Spec}(R)$  be a flat family of stable  $n$ -pointed curves with a single node defined by a section  $Q$  and with the  $n$  smooth points given by sections  $P_\bullet = (P_1, \dots, P_n)$ . Assume that  $\mathcal{C}_0 \setminus P_\bullet(S_0)$  is affine over  $S_0$ . Up to an étale base change of  $S_0$  of degree two, we can normalize  $\mathcal{C}_0$  and obtain a smooth family of  $(n+2)$ -pointed curves  $\tilde{\mathcal{C}}_0 \rightarrow S_0$  with sections  $P_\bullet \sqcup (Q_+, Q_-)$ , where  $Q_\pm(S_0) \in \tilde{\mathcal{C}}_0$  are the preimages of the node in  $\mathcal{C}_0/S_0$ . Fix formal coordinates  $s_+$  and  $s_-$  at  $Q_+(S_0)$  and  $Q_-(S_0)$ , respectively. Such coordinates determine a *smoothing* of  $(\mathcal{C}_0, P_\bullet)$  over  $S = \text{Spec}(R[[q]])$ . That is, a flat family  $\mathcal{C} \rightarrow S = \text{Spec}(R[[q]])$  with sections  $P_\bullet = (P_1, \dots, P_n)$  such that the general fiber is smooth and the special fiber is identified with  $\mathcal{C}_0 \rightarrow S_0$ . The family  $\tilde{\mathcal{C}}_0 \rightarrow S_0$  extends to a family of smooth curves  $\tilde{\mathcal{C}} \rightarrow S = \text{Spec}(R[[q]])$ , with  $n+2$  sections  $P_\bullet, Q_+$ , and  $Q_-$ , with the special fiber identified with  $\tilde{\mathcal{C}}_0 \rightarrow S_0$ , and which fits in the diagram

$$\begin{array}{ccc}
 \tilde{\mathcal{C}} & \xrightarrow{\quad} & \mathcal{C} \\
 \swarrow & & \searrow \\
 & S = \text{Spec}(R[[q]]) & \\
 \nwarrow & & \nearrow \\
 P_\bullet, Q_+, Q_- & & P_\bullet
 \end{array}$$

The formal coordinate at  $Q_\pm(S_0)$  extends to a formal coordinate, still denoted  $s_\pm$ , at  $Q_\pm(S)$  — that is,  $s_\pm$  is generator of the ideal of the completed local  $R[[q]]$ -algebra of  $\tilde{\mathcal{C}}$  at  $Q_\pm(S)$  — such that  $s_+s_- = q$ . For more details, see [Loo, p. 457] and [ACG, pp. 184-5]. We emphasize that the existence of such families holds over the formal base  $S = \text{Spec}(R[[q]])$ , or equivalently, over the complex open unit disk around  $S_0$  in the analytic category, but fails over a more general base. Moreover, one still has that  $\tilde{\mathcal{C}} \setminus P_\bullet(S)$  and  $\mathcal{C} \setminus P_\bullet(S)$  are affine over  $S$ .

**8.2. The sheaf of coinvariants.** The construction of the chiral Lie algebra in §3 and of coinvariants in §4 can be extended over an arbitrary smooth base. One thus obtains a sheaf of Lie algebras  $\mathcal{L}_{\mathcal{C} \setminus P_\bullet}(V)$  and a sheaf of coinvariants  $(\otimes_{i=1}^n M^i \otimes \mathcal{O}_S)_{\mathcal{L}_{\mathcal{C} \setminus P_\bullet}(V)}$ , for given  $V$ -modules  $M^1, \dots, M^n$  and a choice of formal coordinates  $t_i$  at the points  $P_i$ . One can remove the assumption that  $\mathcal{C} \setminus P_\bullet(S)$  is affine over  $S$  using propagation of vacua as

in §4, and furthermore obtain a *sheaf of coinvariants*  $\mathbb{V}(V; M^\bullet)_{(\mathcal{C}/S, P_\bullet)}$  on  $S$  independent of the formal coordinates  $t_i$  via a descent along a torsor as in [FBZ, §17], [DGT2].

Similarly to Proposition 5.1.1, we obtain the following result.

**Theorem 8.2.1.** *Let  $V = \bigoplus_{i \geq 0} V_i$  be a  $C_2$ -cofinite conformal vertex algebra with  $V_0 \cong \mathbb{C}$ . For any collection of finitely generated  $V$ -modules  $M^1, \dots, M^n$ , the sheaf of coinvariants  $\mathbb{V}(V; M^\bullet)_{(\mathcal{C}/S, P_\bullet)}$  is a coherent  $\mathcal{O}_S$ -module.*

*Proof.* The argument runs similarly to the one for Proposition 5.1.1, with the elements of  $\mathcal{L}_{\mathcal{C} \setminus P_\bullet}(V)$  interpreted as elements of  $\mathcal{L}_{\tilde{\mathcal{C}} \setminus P_\bullet \sqcup Q_\bullet}(V)$  satisfying appropriate gluing given by the smoothing construction.  $\square$

Moreover, Theorem 1 holds for such families  $\mathcal{C}_0 \rightarrow S_0$  as follows:

**Theorem 8.2.2.** *Let  $V = \bigoplus_{i \geq 0} V_i$  be a rational,  $C_2$ -cofinite conformal vertex algebra with  $V_0 \cong \mathbb{C}$ . The map (34) induces a canonical  $\mathcal{O}_{S_0}$ -module isomorphism*

$$\mathbb{V}(V; M^\bullet)_{(\mathcal{C}_0/S_0, P_\bullet)} \cong \bigoplus_{W \in \mathcal{W}} \mathbb{V}(V; M^\bullet \otimes W \otimes W')_{(\tilde{\mathcal{C}}_0/S_0, P_\bullet \sqcup Q_\bullet)}.$$

**8.3. Sewing.** Given a simple  $V$ -module  $W = \bigoplus_{i \geq 0} W_i$ , define

$$\mathbf{1}^W := \sum_{i \geq 0} \mathbf{1}^{W_i} q^i \in W \otimes W' \otimes \mathbb{C}[[q]],$$

where  $\mathbf{1}^{W_i} := \text{id}_{W_i} \in \text{End}(W_i) \cong W_i \otimes W_i^\vee$ . Consider the map

$$(36) \quad M^\bullet \longrightarrow M^\bullet \otimes W \otimes W' \otimes \mathbb{C}[[q]], \quad u \mapsto \bigoplus_{W \in \mathcal{W}} u \otimes \mathbf{1}^W.$$

The following result extends [NT, Theorem 8.4.6].

**Theorem 8.3.1.** *Let  $V = \bigoplus_{i \geq 0} V_i$  be a rational,  $C_2$ -cofinite conformal vertex algebra with  $V_0 \cong \mathbb{C}$ , and set  $M^\bullet = \bigotimes_{i=1}^n M^i$  for simple  $V$ -modules  $M^i$ . The map (36) induces a canonical  $\mathcal{O}_{S_0}[[q]]$ -module isomorphism  $\Psi$  such that the following diagram commutes*

$$\begin{array}{ccc} \mathbb{V}(V; M^\bullet)_{(\mathcal{C}/S, P_\bullet)} & \xrightarrow{\Psi} & \bigoplus_{W \in \mathcal{W}} \mathbb{V}(V; M^\bullet \otimes W \otimes W')_{(\tilde{\mathcal{C}}_0/S_0, P_\bullet \sqcup Q_\bullet)} \otimes_{\mathcal{O}_{S_0}} \mathcal{O}_{S_0}[[q]] \\ \downarrow & & \downarrow \\ \mathbb{V}(V; M^\bullet)_{(\mathcal{C}_0/S_0, P_\bullet)} & \xrightarrow{\cong} & \bigoplus_{W \in \mathcal{W}} \mathbb{V}(V; M^\bullet \otimes W \otimes W')_{(\tilde{\mathcal{C}}_0/S_0, P_\bullet \sqcup Q_\bullet)}. \end{array}$$

**Remark 8.3.2.** Observe that by Theorems 8.2.2 and 8.3.1, there is a canonical isomorphism

$$\mathbb{V}(V; M^\bullet)_{(\mathcal{C}/S, P_\bullet)} \cong \mathbb{V}(V; M^\bullet)_{(\mathcal{C}_0/S_0, P_\bullet)}[[q]].$$

In particular, this means that to the non-trivial deformation  $\mathcal{C}$  of  $\mathcal{C}_0$ , there corresponds a trivial deformation of the space of conformal blocks.

*Proof of Theorem 8.3.1.* As in the proof of the Factorization Theorem, we can reduce to the case  $\mathcal{C} \setminus P_\bullet$  affine over  $S$ , we fix formal coordinates  $t_i$  at  $P_i(S)$ , for  $i = 1, \dots, n$ , and  $s_\pm$  at  $Q_\pm(S)$ , and show that (36) induces a canonical  $R[[q]]$ -module isomorphism, still denoted  $\Psi$ , such that the following diagram commutes

$$\begin{array}{ccc} (M^\bullet \otimes \mathcal{O}_S)_{\mathcal{L}_{\mathcal{C} \setminus P_\bullet}(V)} & \xrightarrow{\Psi} & \bigoplus_{W \in \mathcal{W}} (M^\bullet \otimes W \otimes W' \otimes \mathcal{O}_{S_0})_{\mathcal{L}_{\tilde{\mathcal{C}}_0 \setminus P_\bullet \sqcup Q_\bullet}(V)} \otimes_{\mathcal{O}_{S_0}} \mathcal{O}_{S_0}[[q]] \\ \downarrow & & \downarrow \\ (M^\bullet \otimes \mathcal{O}_{S_0})_{\mathcal{L}_{\mathcal{C}_0 \setminus P_\bullet}(V)} & \xrightarrow{\cong} & \bigoplus_{W \in \mathcal{W}} (M^\bullet \otimes W \otimes W' \otimes \mathcal{O}_{S_0})_{\mathcal{L}_{\tilde{\mathcal{C}}_0 \setminus P_\bullet \sqcup Q_\bullet}(V)}. \end{array}$$

The construction will be independent of the choice of the formal coordinates  $t_i$  and  $s_\pm$ , hence this will imply the statement.

*Step 1.* We start by showing that (36) induces a well-defined map  $\Psi$  between spaces of coinvariants. For this, it is enough to show that for each  $\sigma \in \mathcal{L}_{\mathcal{C} \setminus P_\bullet}(V)$  and  $W \in \mathcal{W}$ , one has  $\sigma_{P_\bullet}(M^\bullet) \otimes \mathbf{1}^W = \sigma(M^\bullet \otimes \mathbf{1}^W)$ , or equivalently

$$(37) \quad (\sigma_{Q_+} \otimes 1 + 1 \otimes \sigma_{Q_-})(\mathbf{1}^W) = 0.$$

This vanishing follows by using relative stable differentials on  $\mathcal{C}/S$  to describe elements of  $\mathcal{L}_{\mathcal{C} \setminus P_\bullet}(V)$ , in the same spirit of the proof of Proposition 3.3.1. Namely, an element of  $\mathcal{L}_{\mathcal{C} \setminus P_\bullet}(V)$  can be lifted to one in

$$\bigoplus_{k \geq 0} H^0\left(\mathcal{C} \setminus P_\bullet, \left(\omega_{\mathcal{C}/S}^{\otimes 1-k}\right)^{\oplus \dim V_k}\right).$$

Members of this set can in turn be described as elements in

$$\bigoplus_{k \geq 0} H^0\left(\tilde{\mathcal{C}} \setminus P_\bullet \sqcup Q_\bullet, \left(\omega_{\tilde{\mathcal{C}}/S}^{\otimes 1-k}\right)^{\oplus \dim V_k}\right)$$

satisfying certain order and residue conditions. Before expressing these conditions, let us first note that a section  $\mu$  of  $\omega_{\tilde{\mathcal{C}}/S}^{\otimes 1-k}$  on  $\tilde{\mathcal{C}} \setminus P_\bullet$  satisfies

$$\begin{aligned} \mu_{Q_+} &= \sum_{i,j \geq 0} a_{i+k-1,j} s_+^{i-j+k-1} q^j (ds_+)^{1-k}, \\ \mu_{Q_-} &= \sum_{i,j \geq 0} b_{i,j+k-1} s_-^{j-i+k-1} q^i (ds_-)^{1-k} \end{aligned}$$

with

$$a_{i+k-1,j} = (-1)^{k-1} b_{i,j+k-1}, \quad \text{for } i, j, k \geq 0.$$

This description follows from the identities  $s_+ ds_- + s_- ds_+ = 0$ ,  $s_+ s_- = q$ , and the order conditions on relative stable differentials prescribing that the above two sums are only over nonnegative values of  $i$  and  $j$  (see e.g., [NT, §8.5]). The order conditions for relative stable differentials imply analogous order conditions for elements of  $\mathcal{L}_{\mathcal{C} \setminus P_\bullet}(V)$ . Moreover, the gluing conditions

for elements of  $\mathcal{L}_{\mathcal{E}\setminus P_\bullet}(V)$  are induced from the gluing conditions for sections of  $\mathcal{V}_C$ , see §2.6. Namely, by linearity we can reduce to the case  $\sigma_{Q_+} = A \otimes \mu$  in  $V_k \otimes \omega_{\mathcal{E}/S}^{\otimes 1-k} \cong \left(\omega_{\mathcal{E}/S}^{\otimes 1-k}\right)^{\oplus \dim V_k}$  for  $k \geq 0$ . It follows that  $\sigma_{Q_+}$  and  $\sigma_{Q_-}$  act as

$$\begin{aligned}\sigma_{Q_+} &= \sum_{i,j \geq 0} a_{i+k-1,j} A_{[i-j+k-1]} q^j, \\ \sigma_{Q_-} &= \sum_{i,j \geq 0} b_{i,j+k-1} \sum_{l \geq 0} \frac{1}{l!} (L_1^l A)_{[j-i+k-1-l]} q^i \\ &= \sum_{i,j \geq 0} a_{i+k-1,j} \vartheta (A_{[i-j+k-1]}) q^i.\end{aligned}$$

This is as in the end of the proof of Proposition 3.3.1: the isomorphism between  $V \otimes s_{\pm}^{-1} ds_{\pm} \cong \bigoplus_{k \geq 0} V_k \otimes s_{\pm}^{k-1} (ds_{\pm})^{1-k}$  and the stalk of  $\mathcal{L}_{\mathcal{E}\setminus P_\bullet \sqcup Q_\bullet}(V)$  at  $Q_{\pm}$  maps  $L_1^l A \otimes s_{\pm}^{-1} ds_{\pm}$  to  $A_{[k-l-1]}$ , for  $A \in V_k$ . Hence, the vanishing (37) follows from the identity

$$(38) \quad (A_{[i-j+k-1]} \otimes 1 + 1 \otimes \vartheta (A_{[i-j+k-1]}) q^{i-j}) \mathbf{1}^W = 0$$

established in [NT, Lemma 8.7.1] (note that there is a sign of difference between the involution  $\vartheta$  used in this paper and the involution used in [NT]). Thus we conclude that the map  $\Psi$  is well-defined and makes the diagram above commute.

*Step 2.* Since (i) the target of  $\Psi$  is a free  $\mathcal{O}_{S_0}[[q]]$ -module of finite rank, (ii) the source is finitely generated (Theorem 8.2.1), and (iii)  $\Phi$  is an isomorphism modulo  $q$  (Theorem 8.2.2), Nakayama's lemma implies that  $\Psi$  is an isomorphism (this is as in [TUY], [Loo], [NT]).  $\square$

**Remark 8.3.3.** By [DGT2, §7] there is a twisted logarithmic  $\mathcal{D}$ -module structure on sheaves of coinvariants. Specifically, both  $\mathbb{V}(V; M^\bullet)_{(\mathcal{E}/S, P_\bullet)}$  and  $\bigoplus_{W \in \mathcal{W}} \mathbb{V}(M^\bullet \otimes W \otimes W')_{\mathcal{E}_0 \setminus P_\bullet \sqcup Q_\bullet} \otimes_{\mathcal{O}_{S_0}} \mathcal{O}_{S_0}[[q]]$  are equipped with a projective action by those derivations of  $\mathcal{O}_{S_0}[[q]]$  preserving the ideal generated by  $q$ .

From this point of view, one can interpret Theorem 8.3.1 as a spectral decomposition for the action of  $D = q\partial_q$ . Observe that  $\mathcal{T}_S(\log S_0) = \mathcal{O}_S \otimes_{\mathcal{O}_{S_0}} \mathcal{T}_{S_0} + \mathcal{O}_S q\partial_q$ , and by  $\mathcal{O}_{S_0}[[q]]$ -linearity, the action of  $\mathcal{T}_{S_0}$  commutes with  $\Psi$  since it acts on both spaces by coefficient-wise derivation.

We claim that for every  $u \in M^\bullet$  and  $W \in \mathcal{W}$ , one has

$$(39) \quad \Psi \circ D = (q\partial_q + c_W \text{id}) \circ \Psi,$$

where  $c_W$  is the conformal dimension of  $W$ :  $L_0(w) = (c_W + \deg w)w$ , for homogeneous  $w \in W$ . As in [Loo, Lemma 33], the smoothing construction gives a natural way to lift  $q\partial_q$  to a derivation of  $(\mathcal{O}_{S_0}[[s_+]] \oplus \mathcal{O}_{S_0}[[s_-]])[[q]]$  of the form

$$D_{\underline{a}} := (s_+ \partial_{s_+}, 0) + \sum_{m,n \geq 0} a_{m,n} (s_+^{m-n+1} q^n \partial_{s_+}, -s_-^{n-m+1} q^m \partial_{s_-})$$

for some  $a_{m,n} \in \mathcal{O}_{S_0}$  (there is a typo in [Loo, Lemma 33], nevertheless the computation there is compatible with the above statement). As a special case of the identity (38), we observe that

$$(L_{m-n}q^n, -L_{n-m}q^m)\mathbf{1}^W = 0.$$

It follows that

$$\begin{aligned} q\partial_q(\Psi(u)) &= D_{\underline{a}}(u \otimes \mathbf{1}^W) = D_{\underline{a}}(u) \otimes \mathbf{1}^W + u \otimes D_{\underline{a}}(\mathbf{1}^W) \\ &= D(u) \otimes \mathbf{1}^W + u \otimes (-L_0, 0)(\mathbf{1}^W) \\ &= D(u) \otimes \mathbf{1}^W - c_W u \otimes \mathbf{1}^W \\ &= \Psi(D(u) - c_W(u)), \end{aligned}$$

hence the claim.

**8.4. Proof of Theorem 2.** By means of Theorems 8.2.1 and 8.3.1, one concludes that the sheaf  $\mathbb{V}(V; M^\bullet)$  is a vector bundle of finite rank on  $\overline{\mathcal{M}}_{g,n}$ , as in [TUY], [Sor, §2.7], [Loo], [NT]. Below we sketch the argument for completeness.

*Proof of Theorem 2.* The sheaf  $\mathbb{V}(V; M^\bullet)$  on  $\mathcal{M}_{g,n}$  has finite-dimensional fibers (Proposition 5.1.1) and is equipped with a projectively flat connection [DGT2]. As in [TUY], see also [Sor, §2.7], it follows that  $\mathbb{V}(V; M^\bullet)$  is coherent and locally free of finite rank on  $\mathcal{M}_{g,n}$ . After Theorem 8.2.1 and gluing the sheaf as in [BL], it follows that the sheaf  $\mathbb{V}(V; M^\bullet)$  is also coherent on  $\overline{\mathcal{M}}_{g,n}$ . It remains to show that  $\mathbb{V}(V; M^\bullet)$  is locally free on  $\overline{\mathcal{M}}_{g,n}$ . For this, consider a family of nodal curves  $(\mathcal{C}_0 \rightarrow \text{Spec}(R), P_\bullet)$ , and for simplicity assume that it has only one node. Consider its formal smoothing  $(\mathcal{C} \rightarrow \text{Spec}(R[[q]]), P_\bullet)$  as described in §8.1. Theorem 8.3.1 implies that  $\mathbb{V}(V; M^\bullet)_{(\mathcal{C}/S, P_\bullet)}$  is locally free of finite rank, hence we conclude the argument. For families of curves with more nodes, one proceeds similarly by induction on the number of nodes.  $\square$

## 9. EXAMPLES

In Theorems 1 and 2 and the results leading up to them, including the Sewing Theorem 8.3.1, we assume that the conformal vertex algebras we consider satisfy three hypotheses: (1)  $V = \bigoplus_{i \in \mathbb{Z}_{\geq 0}} V_i$  with  $V_0 \cong \mathbb{C}$ ; (2)  $V$  is rational; and (3)  $V$  is  $C_2$ -cofinite. Here we briefly discuss examples where these hypotheses are known to hold.

**9.1. Virasoro vertex algebras.** Recall the Virasoro Lie algebra  $\text{Vir}$  described in §1.1.1, and consider its Lie subalgebra  $\text{Vir}_{\geq 0} := \mathbb{C}K \oplus z\mathbb{C}[[z]]\partial_z$ . Given complex numbers  $c$  and  $h$ , the Verma module for  $\text{Vir}$  of central charge  $c$  and highest weight  $h$  is  $M_{c,h} := U(\text{Vir}) \otimes_{U(\text{Vir}_{\geq 0})} \mathbb{C}\mathbf{1}$ , where  $\mathbb{C}\mathbf{1}$  is given the structure of a  $\text{Vir}_{\geq 0}$ -module by setting  $L_{p>0}\mathbf{1} = 0$ ,  $L_0\mathbf{1} = h\mathbf{1}$ , and  $K\mathbf{1} = c\mathbf{1}$ . There is a unique maximal proper submodule  $J_{c,h} \subset M_{c,h}$ . For  $h = 0$ ,  $J_{c,0}$

contains a submodule generated by the singular vector  $L_{-1}\mathbf{1} \in M_{c,0}$  [FF]. Set

$$L_{c,h} := M_{c,h}/J_{c,h}, \quad M_c := M_{c,0}/\langle L_{-1}\mathbf{1} \rangle, \quad \text{and} \quad \text{Vir}_c := L_{c,0}.$$

If  $c \neq c_{p,q} := 1 - \frac{6(p-q)^2}{pq}$ , with relatively prime  $p, q \in \mathbb{N}$  such that  $1 < p < q$ , then  $M_c \cong \text{Vir}_c$ , that is,  $J_{c,0} = \langle L_{-1}\mathbf{1} \rangle$ , while for  $c = c_{p,q}$ , the submodule  $J_{c,0}$  is generated by two singular vectors [FF]. By [FZ, Thm 4.3],  $M_c$  and  $\text{Vir}_c$  are conformal vertex algebras and satisfy property (1). By [FZ, Thm 4.6], the Zhu algebra satisfies  $A(M_c) \cong \mathbb{C}[x]$ . As  $\mathbb{C}[x]$  has infinitely many 1-dimensional irreducible representations,  $M_c$  is never rational, and  $\text{Vir}_c$  is not rational for  $c \neq c_{p,q}$ . On the other hand, for  $c = c_{p,q}$ ,  $\text{Vir}_c$  is rational [Wan, Thm 4.2]: the Zhu algebra is  $A(\text{Vir}_c) \cong \mathbb{C}[x]/\langle G_{p,q}(x) \rangle$ , for some  $G_{p,q}(x) \in \mathbb{C}[x]$ , and the irreducible representations of  $\text{Vir}_c$  are given by the modules  $L_{c,h}$ , for  $h = \frac{(np-mq)^2 - (p-q)^2}{4pq}$ , with  $0 < m < p$  and  $0 < n < q$ . Moreover,  $\text{Vir}_c$  is  $C_2$ -cofinite for  $c = c_{p,q}$  [DLM3, Lemma 12.3] (see also [Ara1, Prop. 3.4.1]).

**9.2. Irreducible affine vertex algebras from simple Lie algebras.** We briefly describe the irreducible conformal vertex algebra  $L_\ell(\mathfrak{g})$  associated to a finite-dimensional complex simple Lie algebra  $\mathfrak{g}$  and a level  $\ell \in \mathbb{Z}_{>0}$  [FZ, §2], [LL, §6.2] (see also [NT, §A.1.1]).

Let  $\mathfrak{h} \subset \mathfrak{g}$  be the Cartan subalgebra, and  $\theta$  the longest of some choice of positive roots. Normalize the Cartan-Killing form  $(\cdot, \cdot) : \mathfrak{g} \times \mathfrak{g} \rightarrow \mathbb{C}$  so that  $(\theta, \theta) = 2$ . Let  $\widehat{\mathfrak{g}} := \mathbb{C}K \oplus \mathfrak{g} \otimes \mathbb{C}((t))$  be the corresponding affine Lie algebra with  $K$  central and bracket

$$[A \otimes f(t), B \otimes g(t)] = [A, B] \otimes f(t)g(t) - (A, B) (\text{Res}_{t=0} f dg) K.$$

For  $W$  a  $\mathfrak{g}$ -module, we consider the Verma  $\widehat{\mathfrak{g}}$ -module  $W_\ell = U(\widehat{\mathfrak{g}}) \otimes_{U(\widehat{\mathfrak{g}}_{\geq 0})} W$ , where  $W$  is given the structure of a module over  $\widehat{\mathfrak{g}}_{\geq 0} := \mathbb{C}K \oplus \mathfrak{g} \otimes \mathbb{C}[[t]]$  by letting  $\mathfrak{g} \otimes t\mathbb{C}[[t]]$  act by zero and  $K$  by  $\ell \cdot \text{id}_W$ . For instance, let  $W_\lambda$  be the irreducible  $\mathfrak{g}$ -module associated to a weight  $\lambda \in \mathfrak{h}^*$  and write  $W_{\lambda,\ell} = (W_\lambda)_\ell$ . Let  $J_{\lambda,\ell} \subset W_{\lambda,\ell}$  be the unique maximal proper submodule, and let  $h^\vee$  be the dual Coxeter number of  $\mathfrak{g}$ . By [FZ, Thm 2.4.1], for  $\ell \neq -h^\vee$ ,  $W_{0,\ell}$  and the quotient  $L_\ell(\mathfrak{g}) := W_{0,\ell}/J_{0,\ell}$  are conformal vertex algebras satisfying hypothesis (1), and  $L_\ell(\mathfrak{g})$  is irreducible. Frenkel and Zhu show in [FZ, Thm 3.1.3] that  $L_\ell(\mathfrak{g})$  is rational: the Zhu algebra  $A(L_\ell(\mathfrak{g})) = U(\mathfrak{g})/\langle e_\theta^{\ell+1} \rangle$  is a quotient of the universal enveloping algebra for  $\mathfrak{g}$  (here  $e_\theta$  is a nonzero element of the one-dimensional root space  $\mathfrak{g}_\theta$ ), which is semisimple with irreducible  $U(\mathfrak{g})/\langle e_\theta^{\ell+1} \rangle$ -modules equal to the finite-dimensional irreducible  $\mathfrak{g}$ -modules with highest weight  $\lambda$  such that  $(\lambda, \theta) \leq \ell$ , a finite set [Kac]. Consequently, the irreducible  $L_\ell(\mathfrak{g})$ -modules are  $L_{\lambda,\ell} := W_{\lambda,\ell}/J_{\lambda,\ell}$ , for  $\lambda \in \mathfrak{h}^*$ ,  $(\lambda, \theta) \leq \ell$ . That  $L_\ell(\mathfrak{g})$  is  $C_2$ -cofinite follows from [Zhu2] (see also [DLM3, Prop. 12.6], [Ara1, Proposition 3.5.1]). We emphasize that Verma modules associated to simple  $A(L_\ell(\mathfrak{g}))$ -modules are simple. This follows from two facts: (a)  $L_\ell(\mathfrak{g})$ -modules are integrable  $\widehat{\mathfrak{g}}$ -modules from the Bernstein-Gelfand-Gelfand

category  $\mathcal{O}$  [TK]; and (b) the complete reducibility for integrable  $\widehat{\mathfrak{g}}$ -modules from the category  $\mathcal{O}$  [Kac, Thm 10.7]. Note how complete reducibility can fail outside the BGG category  $\mathcal{O}$  for modules of affine Lie algebras. Indeed we know that Verma modules of affine Lie algebras are indecomposable but not irreducible. On the other hand, modules for irreducible affine vertex algebras coincides with modules from the category  $\mathcal{O}$ , hence Verma modules of irreducible affine vertex algebras are irreducible.

**9.3. The moonshine module  $V^\natural$ .** A rational conformal vertex algebra  $V$  is called *holomorphic* if  $V$  is the unique irreducible  $V$ -module. If properties (1) and (3) also hold, then  $V$  must have central charge divisible by 8 [DM1]. One example of such a conformal vertex algebra is the moonshine module  $V^\natural$ , of central charge 24, relevant for a number of reasons, including the fact that  $\text{Aut}(V^\natural)$  is the monster sporadic group [FLM1].

**9.4. Even lattice vertex algebras.** Let  $L$  be a positive-definite even lattice of finite rank, that is, a free abelian group of finite rank, together with a positive definite bilinear form  $(\cdot, \cdot)$  for which  $(\alpha, \alpha) \in 2\mathbb{Z}$  for any  $\alpha \in L$ . Set  $\mathfrak{h} = \mathbb{C} \otimes_{\mathbb{Z}} L$  (which inherits the bilinear form from  $L$ ). Consider the Heisenberg Lie algebra  $\widehat{\mathfrak{h}}$  associated to  $\mathfrak{h}$  and  $(\cdot, \cdot)$ , defined as the one-dimensional central extension of  $\mathfrak{h}((t))$ :

$$0 \rightarrow \mathbb{C}\mathbf{1} \rightarrow \widehat{\mathfrak{h}} \rightarrow \mathfrak{h}((t)) \rightarrow 0,$$

with bracket given by  $[A \otimes f(t), B \otimes g(t)] = -(A, B)\text{Res}f(t)g'(t)dt \cdot \mathbf{1}$ . Let  $\widetilde{\mathcal{H}}$  be the quotient of the completion of the universal enveloping algebra  $U(\widehat{\mathfrak{h}})$  by the ideal generated by the element  $\mathbf{1} - 1$  (here  $\mathbf{1}$  is the central element of  $\widehat{\mathfrak{h}}$  and  $1$  is the unit of the associative algebra  $U(\widehat{\mathfrak{h}})$ ). The algebra  $\widetilde{\mathcal{H}}$  has topological generators  $h_n$ , for  $h \in \mathfrak{h}$  and  $n \in \mathbb{Z}$ . Given  $\lambda \in \mathfrak{h}$  we consider the (infinitely many) Fock representations of  $\widetilde{\mathcal{H}}$

$$\pi_\lambda = \widetilde{\mathcal{H}} \otimes_{\widetilde{\mathcal{H}}_{\geq 0}} \mathbb{C}v_\lambda,$$

where  $\widetilde{\mathcal{H}}_{\geq 0}$  has topological generators  $h_n$  where  $h \in \mathfrak{h}$  and  $n \in \mathbb{N}$ . To define the structure of an  $\widetilde{\mathcal{H}}_{\geq 0}$ -module on the one-dimensional vector space  $\mathbb{C}v_\lambda$ , for  $h \in \mathfrak{h}$ , one sets  $h_n v_\lambda = 0$  for  $n > 0$ , and  $h_0 v_\lambda = (\lambda, h)v_\lambda$ . There is a conformal vertex algebra structure on  $V_L = \bigoplus_{\lambda \in L} \pi_\lambda$ , and the lattice vertex algebra  $V_L$  satisfies property (1) [Bor], [FLM1]. As a vector space, one has a canonical isomorphism  $V_L \cong \pi_0 \otimes \mathbb{C}[L]$ . Here  $\mathbb{C}[L]$  is the group algebra associated to  $L$  (elements are  $e^\lambda$ , for  $\lambda \in L \setminus \{0\}$ ). By [Don],  $V_L$  is rational, and the complete set of  $V_L$ -modules is given by  $\{V_{L+\lambda} : \lambda \in L'/L\}$ , where  $L' = \{\lambda \in L \otimes_{\mathbb{Z}} \mathbb{Q} : (\lambda, \mu) \in \mathbb{Z}, \forall \mu \in L\}$  is the dual lattice. In [DLM1, Thm 3.4] the Zhu algebra  $A(V_L)$  is described explicitly and in [DLM1, Prop. 2.10] the simple  $A(V_L)$ -modules are constructed, giving an alternative proof of rationality. By [DLM3] the  $V_L$  are  $C_2$ -cofinite.

**9.5. Exceptional  $W$ -algebras.** Arakawa [Ara3] shows that a large number of simple  $W$ -algebras, including all the minimal series principal  $W$ -algebras [FKW] and the exceptional  $W$ -algebras of Kac-Wakimoto [KW], are  $C_2$ -cofinite. He proves that the minimal series principal  $W$ -algebras and a large subclass of exceptional affine  $W$ -algebras are rational [Ara2], [Ara4], [AvE] and satisfy property (1).

**9.6. Orbifold vertex algebras.** Given a vertex algebra  $V$  and a group of automorphisms  $G$  of  $V$ , the orbifold vertex algebra  $V^G$  is the  $G$ -invariant vertex subalgebra of  $V$ . Conjecturally, if  $V$  is  $C_2$ -cofinite (and rational) and  $G$  is finite, then  $V^G$  will also be  $C_2$ -cofinite (and rational). Both of these are known to be true in case  $G$  is a finite and solvable group, when  $V$  is a simple conformal vertex algebra satisfying condition (1) [Miy, CM]. In this case,  $V$  and  $V^G$  satisfy (1), (2), and (3).

**9.7. Commutants.** To a vertex algebra  $V$  and a vertex subalgebra  $\mathcal{A} \subset V$ , Frenkel and Zhu [FZ] defined a vertex algebra  $Com(\mathcal{A}, V)$  arising as a vertex subalgebra that commutes with  $\mathcal{A}$ . Conjecturally, if  $\mathcal{A}$  and  $V$  are both  $C_2$ -cofinite and rational, then  $Com(\mathcal{A}, V)$  will be as well. This was proved for the case of parafermion conformal vertex algebras [DR], after prior work [ALY, DLWY, DLY, DW1, DW2, DW3]. These conformal vertex algebras satisfy (1) [DR].

**9.8. Tensor products.** Let  $V^1, \dots, V^j$  be vertex algebras. Then  $V = \bigotimes_{i=1}^j V^i$  is a vertex algebra and by [DMZ],  $V$  is rational if and only if  $V^i$  is rational for all  $i$ . The tensor product of regular (see §B) conformal vertex algebras is regular [DLM2].

#### APPENDIX A. ZHU'S LIE ALGEBRA AND ISOMORPHIC COINVARIANTS

For a smooth curve  $C$  and a quasi-primary generated vertex algebra  $V$  with  $V_0 \cong \mathbb{C}$ , in addition to the chiral Lie algebra  $\mathcal{L}_{C \setminus P_\bullet}(V)$  (§3.1), one also has Zhu's Lie algebra  $\mathfrak{g}_{C \setminus P_\bullet}(V)$ , reviewed in §A.1.

In Proposition A.2.1, we show that when defined,  $\mathfrak{g}_{C \setminus P_\bullet}(V)$  is isomorphic to the image of  $\mathcal{L}_{C \setminus P_\bullet}(V)$  under the Lie algebra homomorphism  $\varphi_{\mathcal{L}}$  (Proposition A.2.1). Nagatomo and Tsuchiya extend the definition of  $\mathfrak{g}_{C \setminus P_\bullet}(V)$  to stable pointed rational curves [NT], and they indicate that their coinvariants are equivalent to those studied by Beilinson and Drinfeld in [BD], suggesting they knew that Proposition A.2.1 holds in that case.

A quasi-primary vector is an element  $A \in V$  such that  $L_1 A = 0$ , and  $V$  is quasi-primary generated if and only if  $L_1 V_1 = 0$  [DLM4]. A vertex algebra  $V = \bigoplus_{i \geq 0} V_i$  with  $V_0 \cong \mathbb{C}$  satisfies  $L_1 V_1 = 0$  if and only if  $V \cong V'$  (see [FHL, §5.3] and [DM2, §2]). In particular, in the results of Huang [Hua3] and Codogni [Cod], the vertex algebras studied are quasi-primary generated.

**A.1. The Lie algebra  $\mathfrak{g}_{C \setminus P_\bullet}(V)$ .** In [Zhu2], given a smooth pointed curve  $(C, P_\bullet)$  and a quasi-primary generated conformal vertex algebra  $V$  for which  $V_0 \cong \mathbb{C}$ , Zhu defines a Lie algebra  $\mathfrak{g}_{C \setminus P_\bullet}(V)$ , generalizing the construction of Tsuchiya, Ueno, and Yamada for affine Lie algebras. Namely, consider

$$(40) \quad \mathfrak{g}_{C \setminus P_\bullet}(V) := \varphi_{\mathfrak{g}} \left( \bigoplus_{k \geq 0} V_k \otimes H^0 \left( C \setminus P_\bullet, \omega_C^{\otimes 1-k} \right) \right)$$

where

$$(41) \quad \varphi_{\mathfrak{g}}: \bigoplus_{k \geq 0} V_k \otimes H^0 \left( C \setminus P_\bullet, \omega_C^{\otimes 1-k} \right) \rightarrow \bigoplus_{i=1}^n \mathfrak{L}_{t_i}(V)$$

is the map induced by

$$B \otimes \mu \mapsto \left( \text{Res}_{t_i=0} Y[B, t_i] \mu_{P_i} (dt_i)^k \right)_{i=1, \dots, n}.$$

Here  $t_i$  is a formal coordinate at the point  $P_i$ ,  $Y[B, t_i] := \sum_{k \in \mathbb{Z}} B_{[k]} t_i^{-k-1}$ , and  $\mu_{P_i}$  is the Laurent series expansion of  $\mu$  at  $P_i$ , the image of  $\mu$  via

$$H^0 \left( C \setminus P_\bullet, \omega_C^{\otimes 1-k} \right) \rightarrow H^0 \left( D_{P_i}^\times, \omega_C^{\otimes 1-k} \right) \simeq_{t_i} \mathbb{C}((t_i))(dt_i)^{1-k}.$$

When  $V$  is assumed to be quasi-primary generated with  $V_0 \cong \mathbb{C}$ , Zhu shows that  $\mathfrak{g}_{C \setminus P_\bullet}(V)$  is a Lie subalgebra of  $\mathfrak{L}(V)^{\oplus n}$ . The argument uses that any *fixed* smooth algebraic curve admits an atlas such that all transition functions are Möbius transformations. Families of stable rational curves also satisfy this property, a fact used in [NT] to extend Zhu's construction to moduli of stable rational curves. Transition functions between charts on families of curves of arbitrary genus are more general, hence the need to consider the more involved construction for the chiral Lie algebra based on the  $(\text{Aut } \mathcal{O})$ -twist of  $V$  in §2.

**A.2. Isomorphism of coinvariants.** When  $\mathfrak{g}_{C \setminus P_\bullet}(V)$  is well-defined and  $C \setminus P_\bullet$  is affine, one can define the space of coinvariants  $M_{\mathfrak{g}_{C \setminus P_\bullet}(V)}^\bullet$  as the quotient of the  $\mathfrak{L}(V)^{\oplus n}$ -module  $M^\bullet$  by the action of the Lie subalgebra  $\mathfrak{g}_{C \setminus P_\bullet}(V)$  of  $\mathfrak{L}(V)^{\oplus n}$ . These spaces were introduced in [Zhu1], and studied also in [AN, NT]. Recall the homomorphisms  $\varphi_{\mathcal{L}}$  from (14) and  $\varphi_{\mathfrak{g}}$  from (41).

**Proposition A.2.1.** *When  $\mathfrak{g}_{C \setminus P_\bullet}(V)$  is well-defined (§A.1), one has*

$$\text{Im}(\varphi_{\mathcal{L}}) \cong \text{Im}(\varphi_{\mathfrak{g}}).$$

*It follows that there exists an isomorphism of vector spaces*

$$M_{\mathfrak{g}_{C \setminus P_\bullet}(V)}^\bullet \cong M_{\mathcal{L}_{C \setminus P_\bullet}(V)}^\bullet.$$

*Proof.* One has

$$(42) \quad \begin{aligned} \bigoplus_{k \geq 0} V_k \otimes H^0 \left( C \setminus P_\bullet, \omega_C^{\otimes 1-k} \right) &\cong H^0 \left( C \setminus P_\bullet, \bigoplus_{k \geq 0} V_k \otimes \omega_C^{\otimes 1-k} \right) \\ &\cong H^0 \left( C \setminus P_\bullet, \bigoplus_{k \geq 0} \left( \omega_C^{\otimes 1-k} \right)^{\oplus \dim V_k} \right). \end{aligned}$$

From Lemma 2.7.1, one has  $\text{gr}_\bullet \mathcal{Y}_C \cong \bigoplus_{k \geq 0} \left( \omega_C^{\otimes -k} \right)^{\oplus \dim V_k}$ . It follows that

$$H^0 \left( C \setminus P_\bullet, \bigoplus_{k \geq 0} \left( \omega_C^{\otimes 1-k} \right)^{\oplus \dim V_k} \right) \cong H^0 \left( C \setminus P_\bullet, \text{gr}_\bullet \mathcal{Y}_C \otimes \omega_C \right).$$

Now by Lemma 2.7.2,

$$(43) \quad H^0(C \setminus P_\bullet, \text{gr}_\bullet \mathcal{V}_C \otimes \omega_C) \cong H^0(C \setminus P_\bullet, \mathcal{V}_C \otimes \omega_C).$$

On the other hand, as  $C \setminus P_\bullet$  is assumed to be affine, one has

$$\mathcal{L}_{C \setminus P_\bullet}(V) \cong H^0(C \setminus P_\bullet, \mathcal{V}_C \otimes \omega_C) / \nabla H^0(C \setminus P_\bullet, \mathcal{V}_C).$$

The map  $\varphi_{\mathcal{L}}$  is induced from the composition

$$(44) \quad H^0(C \setminus P_\bullet, \mathcal{V}_C \otimes \omega_C) \rightarrow \bigoplus_{i=1}^n H^0(D_{P_i}^\times, \mathcal{V}_C \otimes \omega_C) \rightarrow \bigoplus_{i=1}^n \mathfrak{L}_{t_i}(V).$$

The first map is canonical and obtained by restricting sections; the second is (13). By [FBZ, §6.6.9], sections in  $\nabla H^0(D_{P_i}^\times, \mathcal{V}_C)$  act trivially. Hence (44) induces a map from the Lie algebra  $\mathcal{L}_{C \setminus P_\bullet}(V)$  to  $\bigoplus_{i=1}^n \mathfrak{L}_{t_i}(V)$ . It follows that the image of  $\varphi_{\mathcal{L}}$  coincides with the image of  $H^0(C \setminus P_\bullet, \mathcal{V}_C \otimes \omega_C)$  in  $\bigoplus_{i=1}^n \mathfrak{L}_{t_i}(V)$  via (44). Composing (42) and (43), and by the definition of  $\varphi_{\mathfrak{g}}$  in (41), the image of the map in (44) coincides with the image of  $\varphi_{\mathfrak{g}}$ .  $\square$

## APPENDIX B. MODULES AND REGULARITY

We review here the notion of weak/admissible/ordinary modules and regularity, to compare some of the definitions used here with similar notions in the literature.

A *weak*  $V$ -module is a module as defined in §1.2, without any assumption on the grading. An *ordinary*  $V$ -module is a weak  $V$ -module which carries a  $\mathbb{C}$ -grading  $M = \bigoplus_{\lambda \in \mathbb{C}} M_\lambda$  such that  $L_0|_{M_\lambda} = \lambda \text{id}_{M_\lambda}$ ;  $\dim M_\lambda < \infty$ ; and  $M_{\lambda+n} = 0$ , for fixed  $\lambda$  and  $n \ll 0$ . An *admissible* module is a weak  $V$ -module which carries a  $\mathbb{Z}_{\geq 0}$ -grading and satisfies (2).

An ordinary  $V$ -module is admissible, and when  $V$  is rational, simple admissible  $V$ -modules are ordinary [DLM2, Rmk 2.4]. So for rational  $V$ , the simple admissible  $V$ -modules are the same as the simple ordinary modules.

A vertex algebra is *regular* if any weak module is a direct sum of irreducible ordinary modules [DLM2]. For a conformal vertex algebra  $V = \bigoplus_{i \geq 0} V_i$  with  $V_0 \cong \mathbb{C}$ , it has been proved that  $V$  is regular if it is rational and  $\bar{C}_2$ -cofinite [DLM2, Rmk 3.2], [Li3], [ABD, Thm 4.5].

## ACKNOWLEDGEMENTS

The authors are grateful to Jim Lepowsky, Yi-Zhi Huang, and Bin Gui for helpful discussions. Thanks also to Yi-Zhi Huang and Giulio Codogni for comments on a preliminary version of the manuscript. The authors are indebted to [FBZ] for the treatment of chiral Lie algebras and coinvariants on smooth curves, and to [NT] for the arguments on the factorization and sewing for coinvariants by Zhu's Lie algebra on rational curves. Gibney was supported by NSF DMS-1902237.

## REFERENCES

- [ABD] T. Abe, G. Buhl, and C. Dong. Rationality, regularity, and  $C_2$ -cofiniteness. *Trans. Amer. Math. Soc.*, 356(8):3391–3402, 2004. ↑5, ↑44
- [Abe] T. Abe. A  $\mathbb{Z}_2$ -orbifold model of the symplectic fermionic vertex operator superalgebra. *Math. Z.*, 255(4):755–792, 2007. ↑5
- [ACG] E. Arbarello, M. Cornalba, and P. Griffiths. *Geometry of algebraic curves. Volume II. With a contribution by Joseph Daniel Harris*. Berlin: Springer, 2011. ↑35
- [ADCKP] E. Arbarello, C. De Concini, V. G. Kac, and C. Procesi. Moduli spaces of curves and representation theory. *Comm. Math. Phys.*, 117(1):1–36, 1988. ↑12, ↑13
- [ALY] T. Arakawa, C. H. Lam, and H. Yamada. Zhu’s algebra,  $C_2$ -algebra and  $C_2$ -cofiniteness of parafermion vertex operator algebras. *Adv. Math.*, 264:261–295, 2014. ↑42
- [AN] T. Abe and K. Nagatomo. Finiteness of conformal blocks over compact Riemann surfaces. *Osaka J. Math.*, 40(2):375–391, 2003. ↑2, ↑3, ↑5, ↑23, ↑24, ↑25, ↑43
- [Ara1] T. Arakawa. A remark on the  $C_2$ -cofiniteness condition on vertex algebras. *Math. Z.*, 270(1-2):559–575, 2012. ↑23, ↑40, ↑41
- [Ara2] T. Arakawa. Rationality of Bershadsky-Polyakov vertex algebras. *Communications in Mathematical Physics*, 323(2):627–633, 2013. ↑42
- [Ara3] T. Arakawa. Associated varieties of modules over Kac-Moody algebras and  $C_2$ -cofiniteness of  $W$ -algebras. *Int. Math. Res. Not. IMRN*, (22):11605–11666, 2015. ↑42
- [Ara4] T. Arakawa. Rationality of  $W$ -algebras: principal nilpotent cases. *Ann. of Math. (2)*, 182(2):565–604, 2015. ↑42
- [AvE] T. Arakawa and J. van Ekeren. Rationality and fusion rules of exceptional  $W$ -algebras. *arXiv preprint arXiv:1905.11473*, 2019. ↑42
- [BCG<sup>+</sup>] M. Bainbridge, D. Chen, Q. Gendron, S. Grushevsky, and M. Möller. Strata of  $k$ -differentials. *Algebr. Geom.*, 6(2):196–233, 2019. ↑11
- [BD] A. Beilinson and V. Drinfeld. *Chiral algebras*, volume 51 of *Am Mathem Soc Colloquium Publications*. Am Mathem Soc, Providence, RI, 2004. ↑2, ↑22, ↑43
- [Bea] A. Beauville. Conformal blocks, fusion rules and the Verlinde formula. In *Proceedings of the Hirzebruch 65 Conference on Algebraic Geometry (Ramat Gan, 1993)*, volume 9 of *Israel Math. Conf. Proc.*, pages 75–96. Bar-Ilan Univ., Ramat Gan, 1996. ↑1
- [BFM] A. Beilinson, B. Feigin, and B. Mazur. Introduction to algebraic field theory on curves. *Unpublished*, 1991. ↑3, ↑4, ↑5, ↑22, ↑23
- [BL] A. Beauville and Y. Laszlo. Un lemme de descente. *C. R. Acad. Sci. Paris Sér. I Math.*, 320(3):335–340, 1995. ↑39
- [Bor] R. E. Borcherds. Vertex algebras, Kac-Moody algebras, and the Monster. *Proc. Nat. Acad. Sci. U.S.A.*, 83(10):3068–3071, 1986. ↑10, ↑42
- [Buh] G. Buhl. A spanning set for VOA modules. *J. Algebra*, 254(1):125–151, 2002. ↑23, ↑25
- [CM] S. Carnahan and M. Miyamoto. Regularity of fixed-point vertex operator subalgebras. *arXiv preprint arXiv:1603.05645*, 2016. ↑42
- [Cod] G. Codogni. Vertex algebras and teichmüller modular forms. *arXiv:1901.03079*, pages 1–31, 2019. ↑3, ↑4, ↑43
- [DGT1] C. Damiolini, A. Gibney, and N. Tarasca. Chern classes of vector bundles from vertex algebras on moduli of curves. *In preparation*. ↑4
- [DGT2] C. Damiolini, A. Gibney, and N. Tarasca. Conformal blocks from vertex algebras and their connections on  $\overline{\mathcal{M}}_{g,n}$ . *Submitted*, 2019. ↑1, ↑2, ↑3, ↑4, ↑12, ↑14, ↑17, ↑22, ↑30, ↑36, ↑38, ↑39

- [DL] C. Dong and J. Lepowsky. *Generalized vertex algebras and relative vertex operators*, volume 112 of *Progress in Maths*. Birkhäuser Boston, Inc., Boston, MA, 1993. ↑7
- [DLM1] C. Dong, H. Li, and G. Mason. Certain associative algebras similar to  $U(\mathfrak{sl}_2)$  and Zhu's algebra  $A(V_L)$ . *J. Algebra*, 196(2):532–551, 1997. ↑42
- [DLM2] C. Dong, H. Li, and G. Mason. Regularity of rational vertex operator algebras. *Adv. Math.*, 132(1):148–166, 1997. ↑5, ↑42, ↑44
- [DLM3] C. Dong, H. Li, and G. Mason. Modular-invariance of trace functions in orbifold theory and generalized Moonshine. *Comm. Math. Phys.*, 214(1):1–56, 2000. ↑40, ↑41, ↑42
- [DLM4] C. Dong, Z. Lin, and G. Mason. On vertex operator algebras as  $\mathfrak{sl}_2$ -modules. In *Groups, difference sets, and the Monster (Columbus, OH, 1993)*, volume 4 of *Ohio State Univ. Math. Res. Inst. Publ.*, pages 349–362. de Gruyter, Berlin, 1996. ↑43
- [DLWY] C. Dong, C. H. Lam, Q. Wang, and H. Yamada. The structure of parafermion vertex operator algebras. *J. Algebra*, 323(2):371–381, 2010. ↑42
- [DLY] C. Dong, C. H. Lam, and H. Yamada.  $W$ -algebras related to parafermion algebras. *J. Algebra*, 322(7):2366–2403, 2009. ↑42
- [DM1] C. Dong and G. Mason. Holomorphic vertex operator algebras of small central charge. *Pacific journal of mathematics*, 213(2):253–266, 2004. ↑41
- [DM2] C. Dong and G. Mason. Rational vertex operator algebras and the effective central charge. *Int. Math. Res. Not.*, (56):2989–3008, 2004. ↑43
- [DMZ] C. Dong, G. Mason, and Y. Zhu. Discrete series of the Virasoro algebra and the moonshine module. In *Algebraic groups and their generalizations: quantum and infinite-dimensional methods (University Park, PA, 1991)*, volume 56 of *Proc. Sympos. Pure Math.*, pages 295–316. Amer. Math. Soc., Providence, RI, 1994. ↑42
- [Don] C. Dong. Vertex algebras associated with even lattices. *J. Algebra*, 161(1):245–265, 1993. ↑42
- [DR] C. Dong and L. Ren. Representations of the parafermion vertex operator algebras. *Adv. Math.*, 315:88–101, 2017. ↑42
- [DW1] C. Dong and Q. Wang. The structure of parafermion vertex operator algebras: general case. *Comm. Math. Phys.*, 299(3):783–792, 2010. ↑42
- [DW2] C. Dong and Q. Wang. On  $C_2$ -cofiniteness of parafermion vertex operator algebras. *J. Algebra*, 328:420–431, 2011. ↑42
- [DW3] C. Dong and Q. Wang. Parafermion vertex operator algebras. *Front. Math. China*, 6(4):567–579, 2011. ↑42
- [Fak] N. Fakhruddin. Chern classes of conformal blocks. In *Compact moduli spaces and vector bundles*, volume 564 of *Contemp. Math.*, pages 145–176. Amer. Math. Soc., Providence, RI, 2012. ↑22
- [FBZ] E. Frenkel and D. Ben-Zvi. *Vertex algebras and algebraic curves*, volume 88 of *Mathematical Surveys and Monographs*. Am Mathem Soc, Providence, RI, second edition, 2004. ↑1, ↑2, ↑5, ↑7, ↑8, ↑11, ↑12, ↑14, ↑15, ↑16, ↑17, ↑18, ↑22, ↑23, ↑27, ↑28, ↑29, ↑30, ↑32, ↑36, ↑44, ↑45
- [FF] B. L. Feigin and D. B. Fuks. Verma modules over a Virasoro algebra. *Funktsional. Anal. i Prilozhen.*, 17(3):91–92, 1983. ↑40
- [FHL] I. B. Frenkel, Y.-Z. Huang, and J. Lepowsky. On axiomatic approaches to vertex operator algebras and modules. *Mem. Amer. Math. Soc.*, 104(494):viii+64, 1993. ↑7, ↑10, ↑43
- [FKW] E. Frenkel, V. Kac, and M. Wakimoto. Characters and fusion rules for  $W$ -algebras via quantized Drinfel'd-Sokolov reduction. *Comm. Math. Phys.*, 147(2):295–328, 1992. ↑42

- [FLM1] I. B. Frenkel, J. Lepowsky, and A. Meurman. A natural representation of the Fischer-Griess Monster with the modular function  $J$  as character. *Proc. Nat. Acad. Sci. U.S.A.*, 81(10, , Phys. Sci.):3256–3260, 1984. ↑41, ↑42
- [FLM2] I. Frenkel, J. Lepowsky, and A. Meurman. *Vertex operator algebras and the Monster*, volume 134 of *Pure and Applied Maths*. Academic Press, Inc., Boston, MA, 1988. ↑8
- [FZ] I. B. Frenkel and Y. Zhu. Vertex operator algebras associated to representations of affine and Virasoro algebras. *Duke Math. J.*, 66(1):123–168, 1992. ↑7, ↑9, ↑23, ↑40, ↑42
- [Hua1] Y.-Z. Huang. Differential equations and intertwining operators. *Commun. Contemp. Math.*, 7(3):375–400, 2005. ↑3, ↑4
- [Hua2] Y.-Z. Huang. Differential equations, duality and modular invariance. *Commun. Contemp. Math.*, 7(5):649–706, 2005. ↑3, ↑4
- [Hua3] Y.-Z. Huang. Vertex operator algebras, the Verlinde conjecture, and modular tensor categories. *Proc. Natl. Acad. Sci. USA*, 102(15):5352–5356, 2005. ↑4, ↑43
- [Hua4] Y.-Z. Huang. Rigidity and modularity of vertex tensor categories. *Commun. Contemp. Math.*, 10(suppl. 1):871–911, 2008. ↑3, ↑4
- [Hua5] Y.-Z. Huang. Vertex operator algebras and the Verlinde conjecture. *Commun. Contemp. Math.*, 10(1):103–154, 2008. ↑3, ↑4
- [Kac] V. G. Kac. Infinite-dimensional Lie algebras, and the Dedekind  $\eta$ -function. *Funkcional. Anal. i Priložen.*, 8(1):77–78, 1974. ↑41
- [KW] V. G. Kac and M. Wakimoto. On rationality of  $W$ -algebras. *Transform. Groups*, 13(3-4):671–713, 2008. ↑42
- [KZ] V. G. Knizhnik and A. B. Zamolodchikov. Current algebra and Wess-Zumino model in two dimensions. *Nuclear Phys. B*, 247(1):83–103, 1984. ↑23
- [Li1] H.-S. Li. Local systems of vertex operators, vertex superalgebras and modules. *J. Pure Appl. Algebra*, 109(2):143–195, 1996. ↑7
- [Li2] H. Li. *Representation theory and tensor product theory for vertex operator algebras*. ProQuest LLC, Ann Arbor, MI, 1994. Thesis (Ph.D.)—Rutgers The State University of New Jersey - New Brunswick. ↑9
- [Li3] H. Li. Some finiteness properties of regular vertex operator algebras. *J. Algebra*, 212(2):495–514, 1999. ↑44
- [LL] J. Lepowsky and H. Li. *Introduction to vertex operator algebras and their representations*, volume 227 of *Progress in Maths*. Birkhäuser Boston, Inc., Boston, MA, 2004. ↑7, ↑40
- [Loo] E. Looijenga. From WZW models to modular functors. In *Handbook of moduli. Vol. II*, volume 25 of *Adv. Lect. Math. (ALM)*, pages 427–466. Int. Press, Somerville, MA, 2013. ↑22, ↑35, ↑38, ↑39
- [Miy] M. Miyamoto.  $C_2$ -cofiniteness of cyclic-orbifold models. *Comm. Math. Phys.*, 335(3):1279–1286, 2015. ↑42
- [MMS] S. D. Mathur, S. Mukhi, and A. Sen. On the classification of rational conformal field theories. *Phys. Lett. B*, 213(3):303–308, 1988. ↑23
- [MOP1] A. Marian, D. Oprea, and R. Pandharipande. The first Chern class of the Verlinde bundles. In *String-Math 2012*, volume 90 of *Proc. Sympos. Pure Math.*, pages 87–111. Amer. Math. Soc., Providence, RI, 2015. ↑1
- [MOP<sup>+</sup>2] A. Marian, D. Oprea, R. Pandharipande, A. Pixton, and D. Zvonkine. The Chern character of the Verlinde bundle over  $\overline{\mathcal{M}}_{g,n}$ . *J. Reine Angew. Math.*, 732:147–163, 2017. ↑1
- [MS] G. Moore and N. Seiberg. Classical and quantum conformal field theory. *Comm. Math. Phys.*, 123(2):177–254, 1989. ↑1, ↑4
- [NT] K. Nagatomo and A. Tsuchiya. Conformal field theories associated to regular chiral vertex operator algebras. I. Theories over the projective line. *Duke Math.*

- J.*, 128(3):393–471, 2005. ↑2, ↑3, ↑4, ↑5, ↑7, ↑9, ↑10, ↑23, ↑26, ↑29, ↑33, ↑36, ↑38, ↑39, ↑40, ↑42, ↑43, ↑45
- [Seg1] G. Segal. Two-dimensional conformal field theories and modular functors. In *IXth International Congress on Mathematical Physics (Swansea, 1988)*, pages 22–37. Hilger, Bristol, 1989. ↑1
- [Seg2] G. Segal. The definition of conformal field theory. In *Topology, geometry and quantum field theory*, volume 308 of *London Math. Soc. Lecture Note Ser.*, pages 421–577. Cambridge Univ. Press, Cambridge, 2004. ↑1
- [Sor] C. Sorger. La formule de Verlinde. *Astérisque*, (237):Exp. No. 794, 3, 87–114, 1996. Séminaire Bourbaki, Vol. 1994/95. ↑39
- [Tat] J. Tate. Residues of differentials on curves. *Ann. Sci. École Norm. Sup. (4)*, 1:149–159, 1968. ↑27
- [TK] A. Tsuchiya and Y. Kanie. Vertex operators in the conformal field theory on  $\mathbf{P}^1$  and monodromy representations of the braid group. *Lett. Math. Phys.*, 13(4):303–312, 1987. ↑4, ↑23, ↑41
- [Tsu] Y. Tsuchimoto. On the coordinate-free description of the conformal blocks. *J. Math. Kyoto Univ.*, 33(1):29–49, 1993. ↑4
- [TUY] A. Tsuchiya, K. Ueno, and Y. Yamada. Conformal field theory on universal family of stable curves with gauge symmetries. In *Integrable systems in quantum field theory and statistical mechanics*, volume 19 of *Adv. Stud. Pure Math.*, pages 459–566. Academic Press, Boston, MA, 1989. ↑3, ↑4, ↑5, ↑23, ↑27, ↑38, ↑39
- [Vak] R. Vakil. The rising sea. *Foundations of Algebraic Geometry*. Available at [math216.wordpress.com](http://math216.wordpress.com), 2017. ↑19
- [Wan] W. Wang. Rationality of Virasoro vertex operator algebras. *Internat. Math. Res. Notices*, (7):197–211, 1993. ↑40
- [Zhu1] Y. Zhu. Global vertex operators on Riemann surfaces. *Comm. Math. Phys.*, 165(3):485–531, 1994. ↑1, ↑2, ↑12, ↑23, ↑43
- [Zhu2] Y. Zhu. Modular invariance of characters of vertex operator algebras. *J. Amer. Math. Soc.*, 9(1):237–302, 1996. ↑2, ↑6, ↑7, ↑8, ↑9, ↑23, ↑26, ↑41, ↑43

CHIARA DAMIOLINI

DEPARTMENT OF MATHEMATICS, PRINCETON UNIVERSITY, PRINCETON, NJ 08540

*E-mail address:* [chiarad@princeton.edu](mailto:chiarad@princeton.edu)

ANGELA GIBNEY

DEPARTMENT OF MATHEMATICS, RUTGERS UNIVERSITY, PISCATAWAY, NJ 08854

*E-mail address:* [angela.gibney@rutgers.edu](mailto:angela.gibney@rutgers.edu)

NICOLA TARASCA

DEPARTMENT OF MATHEMATICS & APPLIED MATHEMATICS,  
VIRGINIA COMMONWEALTH UNIVERSITY, RICHMOND, VA 23284

*Previous address:*

DEPARTMENT OF MATHEMATICS, RUTGERS UNIVERSITY, PISCATAWAY, NJ 08854

*E-mail address:* [tarascan@vcu.edu](mailto:tarascan@vcu.edu)