

# Angela Gibney

Moduli spaces reveal how objects like varieties or schemes of a particular type behave in families. For instance, the moduli space  $\overline{\mathcal{M}}_{g,n}$ , parametrizing stable  $n$ -pointed curves of genus  $g$ , gives insight into the study of smooth curves and their degenerations. As curves arise in many contexts, the moduli space of curves is a meeting ground, where constructions from diverse fields can be tested and explored.

A natural goal when studying a space, like  $\overline{\mathcal{M}}_{g,n}$ , is to characterize the maps admitted by it. From this perspective, vector bundles are important, as their sections can be thought of as functions. My recent work concerns vector bundles on  $\overline{\mathcal{M}}_{g,n}$  defined from representations of vertex operator algebras, generalizing Verlinde bundles, defined by integrable modules over affine Lie algebras. In earlier research, I studied aspects the moduli space of curves, using Mori theory and tropical geometry.

Here I briefly describe a selection of my results, arranged according to the following themes:

- (1) Bundles on the moduli space of curves from modules over vertex algebras;
- (2) Verlinde bundles and the geometry of the moduli space of curves;
- (3) Moduli, reductions, and generalizations of the F-Conjecture.

## 1. BUNDLES ON THE MODULI SPACE OF CURVES FROM MODULES OVER VERTEX OPERATOR ALGEBRAS

For years it has been understood that coinvariants defined from the action of a Lie algebra on modules  $W^i$  over a vertex operator algebra satisfying certain properties give rise to sheaves  $\mathbb{V}_g(V; W^\bullet)$  on moduli of *smooth*  $n$ -pointed curves of genus  $g$  [FBZ04]. In the special cases of modules over Virasoro vertex algebras, and of integrable modules over affine Lie algebras, the sheaves were shown to extend to stable curves with singularities, and have particularly nice properties [TK88, TUY89, BDF91]. For instance, they were shown to be locally free of finite rank, giving vector bundles. Those vector bundles satisfy factorization (and the more refined sewing property), a feature that makes recursive arguments about ranks and Chern classes possible. For the latter type, defined from integrable modules over affine Lie algebras, the sheaves are sometimes referred to as Verlinde bundles as their ranks can be computed with the Verlinde formula [MOP15, MOP+17]. The Verlinde bundles support a projectively flat connection, and their Chern characters determine cohomological field theories [TUY89, Tsu93, MOP15, MOP+17].

There have been questions as to whether the more general sheaves of modules over vertex operator algebras extend to stable curves with singularities, and if so, what conditions would ensure they satisfy the valuable properties known to hold for the Verlinde bundles [Zhu94, FBZ04]. Results for  $g = 0$ , and  $g = 1$  gave support to speculation and conjecture regarding what should suffice [AN03a, AN03b, NT05, Hua05a, Hua05b, Hua05c, Hua08].

With my current and former postdocs C. Damiolini and N. Tarasca, I have identified three properties that if satisfied by a vertex operator algebra  $V$ , guarantee that all features mentioned above hold. Namely, if  $V$  is (I) of CFT-type; (II) rational; and (III)  $C_2$ -cofinite, then sheaves of coinvariants defined from finitely generated admissible modules extend to the boundary, are locally free of finite rank, satisfy factorization, support a projectively flat connection, and give way to cohomological field theories:

*Sheaves on  $\overline{\mathcal{M}}_{g,n}$  and their projectively flat connection.* In [DGT19a], we show that coinvariants defined by simple admissible modules  $W^i$  over a vertex operator algebras satisfying properties (I), (II), and (III) define quasi-coherent sheaves  $\mathbb{V}_g(V; W^\bullet)$  on the moduli space  $\overline{\mathcal{M}}_{g,n}$  of *stable*  $n$ -pointed curves of genus  $g$ . We show that these sheaves carry a twisted logarithmic D-module structure, and hence support a projectively flat connection.

*Vector bundles, factorization, and sewing.* In [DGT19b] we show that vector spaces of coinvariants defined by simple admissible modules  $W^i$  over a vertex operator algebra satisfying properties (I), (II), and (III) satisfy factorization, and the sewing theorem, a version of factorization on a smoothing of a nodal curve, solving a 25 year old conjecture. As an application, we prove the sheaves  $\mathbb{V}_g(V; W^\bullet)$  are locally free and of finite rank, giving vector bundles on  $\overline{\mathcal{M}}_{g,n}$ . This has led to the recent solution of longstanding questions about convergence of the sewing map in the analytic setting [Gui20].

*Vertex algebras of CohFT-type.* In [DGT19c] we show that, as for Verlinde bundles, those given by finitely generated admissible modules over a conformal vertex algebra satisfying properties (I), (II), and (III) give rise to cohomological field theories. Following [MOP<sup>+</sup>17], we give a polynomial expression (with rational coefficients) for their Chern characters. In this paper we say that if they satisfy properties (I), (II), and (III), then the vertex algebra is of CohFT-type.

*Current work.* Given the similarities observed between bundles of coinvariants defined by simple admissible modules over a vertex operator algebra of CohFT-type with Verlinde bundles, it is expected that these generalized Verlinde bundles share other important properties with their classical counterparts. For instance, Verlinde bundles were shown to be globally generated on  $\overline{\mathcal{M}}_{0,n}$  by Fakhruddin [Fak12], and we hope to prove global generation of bundles  $\mathbb{V}_0(V; W^\bullet)$ , for vertex operator algebras of CohFT-type if  $V$  is generated in degree 1, or if the rank of  $\mathbb{V}_0(V; W^\bullet)$  is 1. This could lead to new information about the birational geometry of the moduli space of curves. First Chern classes of globally generated bundles are base point free and give rise to morphisms. Verlinde bundles were shown to define maps from  $\overline{\mathcal{M}}_{0,n}$  to moduli spaces parametrizing weighted points supported on (generalized) Veronese curves [Gia13, GG12], generalizing Kapranov's constructions in [Kap93a], [Kap93b].

## 2. VERLINDE BUNDLES AND THE GEOMETRY OF THE MODULI SPACE OF CURVES

As discussed above, coinvariants defined by integrable modules over affine Lie algebras define vector bundles on  $\overline{\mathcal{M}}_{g,n}$ , sometimes called Verlinde bundles as their ranks can be computed with Verlinde formula. These have special properties: Dual spaces to fibers have connections to generalized theta functions, and in genus zero bundles are known to be globally generated, and so their Chern classes have positivity properties. Some of my results about these objects are briefly described below.

*Basepoint free Gromov-Witten loci on  $\overline{\mathcal{M}}_{0,n}$ .* In [BG18], with P. Belkale, we study basepoint free loci on  $\overline{\mathcal{M}}_{0,n}$  defined from the Gromov-Witten invariants of smooth projective homogeneous spaces  $X$ . When  $X$  is projective space, divisors are shown equivalent to first Chern classes of Verlinde bundles defined by integrable modules at level one over simple Lie algebras in type A.

*Current work.* In [BG18], we have conjectured that for  $X = \text{Gr}(r, r + \ell)$ , there is an extension of the level  $\ell = 1$  identity mentioned above, to first Chern classes in type A at the critical level  $\ell$ , giving a higher rank version of Witten's theorem, identifying ranks of Verlinde bundles in type A with iterated structure constants in the quantum cohomology ring of the Grassmannian. Together with Linda Chen, I am currently working with graduate students L Heller (UC Berkeley), H Larson (Stanford), and postdoc E Kalashnikov (Harvard), continuing a project started at the ICERM Women in Algebraic Geometry workshop (online) this summer, studying these classes, and working on this conjecture.

*Finite generation of the determinate of cohomology line bundle.* For  $G$  a simple, simply connected complex linear algebraic group,  $C$  stable of genus  $g \geq 2$ ,  $\text{Bun}_G(C)$  is the stack of principal  $G$ -bundles on  $C$ . To any representation  $G \rightarrow \text{GL}(V)$ , let  $\mathcal{D}(V)$  on  $\text{Bun}_G(C)$ , be the determinant of cohomology line bundle. In [BG], we show, when  $G$  is  $\text{SL}(r)$ , and for the standard representation  $\text{SL}(r) \rightarrow \text{GL}(V)$ , one has that  $\bigoplus_{m \in \mathbb{Z}_{\geq 0}} H^0(\text{Bun}_{\text{SL}(r)}(C), \mathcal{D}(V)^{\otimes m})$  is finitely generated. This was known for smooth curves [BL94, Fal94]. As an application, using Verlinde bundles, we show there is a flat family over  $\overline{\mathcal{M}}_g$  whose fibers are normal and projective, and whose fibers over smooth  $C$  are moduli spaces  $\text{SU}_C(r)$ , parameterizing semistable vector bundles of rank  $r$  with trivial determinant on  $C$ .

*Geometric interpretations at points on the boundary.* At smooth curves, conformal blocks for affine Lie algebras correspond to generalized theta functions [BL94, Fal94, KNR94, Pau96, LS97], and in [BG, Theorems 1.3 and 1.4], with P. Belkale, we show this is true for singular curves as well, in prescribed circumstances. In [BGK16], with P. Belkale and A. Kazanova, we show using intersection theory on  $\overline{\mathcal{M}}_g$ , this is not true in all situations.

*Vanishing and identities.* Characteristic classes of Verlinde bundles are subject to a number of types of identities, which can give information about their associated maps. For instance, in [BGM15], with P. Belkale and S. Mukhopadhyay, we found new rank and level identities satisfied by first Chern classes of bundles defined by integrable modules over affine Lie algebras of type  $A$  at the critical level, generalizing work of Fakhruddin in type  $A_1$ . Inspired by unexpected examples instances of trivial divisors, in [BGM16] we introduce the problem of finding necessary and sufficient conditions to determine when the first Chern class is nontrivial (solving the problem in type  $A_1$ ). We also show under certain hypothesis, first Chern classes are subject to additive identities. Identities for first Chern classes were also found in [AGSS12], and [AGS14], and in [GM16], extended to higher Chern classes.

*Birational models.* Verlinde bundles have helped to provide modular interpretations of images of birational contractions of  $\overline{\mathcal{M}}_{0,n}$ , including GIT quotients generalizing Kapranov's compactifications of  $\mathcal{M}_{0,n}$  [Kap93a, Kap93b], generalized by Giansiracusa and Simpson. In [GG] and [GJMS13] with my coauthors Giansiracusa, and Giansiracusa, Jensen, Moon, and Swinarski, this is further developed.

*Finite generation of full-dimensional subcones of the Nef cone.* In [GG12], with Giansiracusa, we show the infinite set of first Chern classes on  $\overline{\mathcal{M}}_{0,n}$  of bundles defined by integrable modules at level one over affine Lie algebras in type  $A$  spans a finitely generated, full-dimensional subcone of the cone of nef divisors. This was generalized in different contexts to full dimensional cones generated by infinite sets of first Chern classes of rank one bundles by [Kaz16], my former postdoc, and [Hob19], my student.

*Conformal blocks and the Torelli map from  $\overline{\mathcal{M}}_g$  to certain compactifications of  $A_g$ .* While not known to be base point free generally, conformal blocks divisors can sometimes be used to study moduli of higher genus curves. In [Gib12], I identify one of the faces of the nef cone of  $\overline{\mathcal{M}}_g$ , consisting of semi-ample divisors, showing they are pullbacks of ample divisors along the Torelli map from  $\overline{\mathcal{M}}_g$  to certain types of compactifications of  $A_g$ , the moduli space of Abelian varieties of dimension  $g$ .

### 3. GENERALIZATIONS OF THE MODULI OF CURVES, TROPICAL COMPACTIFICATIONS AND THE F-CONJECTURE.

*Pointed trees of projective spaces.* In [CGK09], Chen, Krashen and I construct a space  $T_{d,n}$ , whose points correspond to  $n$ -pointed stable rooted trees of  $d$ -dimensional projective spaces, which for  $d = 1$ , are  $(n + 1)$ -pointed stable rational curves. Smooth and projective, with boundary a smooth normal crossings divisor, the space  $T_{d,n}$  has an inductive construction analogous to but differing from Keel's construction of  $\overline{\mathcal{M}}_{0,n}$ . We describe its Chow groups, Chow ring, Chow motive, and Poincaré polynomials, and use this information to answer a question of Fulton and MacPherson.

*Reduction of the F-Conjecture to genus zero.* The F-Conjecture asserts that a divisor on  $\overline{\mathcal{M}}_{g,n}$  is nef if and only if it nonnegatively intersects one of a finite number of rational curves on  $\overline{\mathcal{M}}_{g,n}$  called F-Curves. In particular, the F-Conjecture implies that  $\text{Nef}(\overline{\mathcal{M}}_{g,n})$  is polyhedral, the convex hull of a finite number of extremal rays.

In [GKM02], we proved that the F-Conjecture on  $\overline{\mathcal{M}}_{0,g+n}$  implies the F-Conjecture on  $\overline{\mathcal{M}}_{g,n}$ . The conjecture holds for low  $g$  and  $n$  [KM13, GKM02, FG03, Gib09].

*Reduction of the F-Conjecture to log canonical.* In [Gib09], I reduce the F-conjecture on  $\overline{\mathcal{M}}_{g,n}$  to showing that certain divisors in  $\overline{\mathcal{M}}_{0,N}$  for  $N \leq g+n$  are equivalent to the sum of the canonical divisor plus an effective divisor supported on the boundary. As an application, I give numerical criteria, which if satisfied by a divisor  $D$  on  $\overline{\mathcal{M}}_g$ , show that  $D$  is nef. Using computer software, written by Krashen, this is used to verify the conjecture for  $g \leq 24$ .

*Generalization to tropical compactifications.* While generally not a toric variety, it is possible to embed  $\overline{\mathcal{M}}_{0,n}$  into a noncomplete toric variety  $X_\Delta$ , and in [GM10], we give equations for  $\overline{\mathcal{M}}_{0,n}$  in the Cox ring of  $X_\Delta$ . In [GM12], we study varieties  $X$  that, like  $\overline{\mathcal{M}}_{0,n}$ , can be embedded in a toric variety  $X_\Delta$ , as a tropical compactification. We use  $X_\Delta$  to define cones which are upper and lower bounds for  $\text{Nef}(X)$ . For  $X = \overline{\mathcal{M}}_{0,n}$ , we show that our upper bound cone  $U(\overline{\mathcal{M}}_{0,n})$ , is the cone of divisors predicted by the F-Conjecture to be equal to  $\text{Nef}(\overline{\mathcal{M}}_{0,n})$ . In other words, this result generalizes the F-Conjecture on  $\overline{\mathcal{M}}_{0,n}$  to varieties that like  $\overline{\mathcal{M}}_{0,n}$ , are tropical compactifications.

### REFERENCES

- [AN03a] Toshiyuki Abe and Kiyokazu Nagatomo, *Finiteness of conformal blocks over the projective line*, Vertex operator algebras in mathematics and physics (Toronto, ON, 2000), Fields Inst. Commun., vol. 39, Amer. Math. Soc., Providence, RI, 2003, pp. 1–12. [↑1](#)
- [AN03b] ———, *Finiteness of conf blocks over compact Riemann surfaces*, Osaka J. Math. **40** (2003), no. 2, 375–391. [↑1](#)
- [AGS14] Valery Alexeev, Angela Gibney, and David Swinarski, *Higher-level  $\mathfrak{sl}_2$  conformal blocks divisors on  $\overline{\mathcal{M}}_{0,n}$* , Proc. Edinb. Math. Soc. (2) **57** (2014), no. 1, 7–30. [↑3](#)
- [AGSS12] Maxim Arap, Angela Gibney, James Stankewicz, and David Swinarski,  *$\mathfrak{sl}_n$  level 1 conformal blocks divisors on  $\overline{\mathcal{M}}_{0,n}$* , Int. Math. Res. Not. IMRN **7** (2012), 1634–1680. [↑3](#)
- [BL94] Arnaud Beauville and Yves Laszlo, *Conformal blocks and generalized theta functions*, Comm. Math. Phys. **164** (1994), no. 2, 385–419. [↑3](#)
- [BG18] Prakash Belkale and Angela Gibney, *Basepoint free cycles on  $\overline{\mathcal{M}}_{0,n}$  from Gromov-Witten theory*, arXiv:1809.02986 [math.AG] (2018). [↑2](#)
- [BG] ———, *On finite generation of the section ring of the determinant of cohomology line bundle*, Transactions of the AMS, to appear. [↑3](#)

- [BGM16] Prakash Belkale, Angela Gibney, and Swarnava Mukhopadhyay, *Nonvanishing of conformal blocks divisors on  $\overline{M}_{0,n}$* , *Transform. Groups* **21** (2016), no. 2, 329–353. [↑3](#)
- [BGM15] ———, *Vanishing and identities of conformal blocks divisors*, *Algebr. Geom.* **2** (2015), no. 1, 62–90. [↑3](#)
- [BGK16] Prakash Belkale, Angela Gibney, and Anna Kazanova, *Scaling of conformal blocks and generalized theta functions over  $\overline{M}_{g,n}$* , *Mathematische Zeit.* (2016), 1–27. [↑3](#)
- [BDF91] A. Beilinson, V. Drinfeld, and Boris Feigin, *Introd to Algebraic Field Theory on Curves*, unpublished (1991). [↑1](#)
- [CGK09] L. Chen, A. Gibney, and D. Krashen, *Pointed trees of projective spaces*, *J. Algebraic Geom.* **18** (2009), no. 3, 477–509. [↑4](#)
- [DGT19a] Chiara Damiolini, Angela Gibney, and Nicola Tarasca, *Conformal blocks from vertex algebras and their connections on  $\overline{M}_{g,n}$*  (2019). *Geometry and Topology*, to appear, <https://arxiv.org/abs/1901.06981>. [↑1](#)
- [DGT19b] ———, *On factorization of conformal blocks from vertex algebras* (2019). <https://arxiv.org/abs/1909.04683>. [↑2](#)
- [DGT19c] ———, *Vertex algebras of CohFT-type* (2019). to appear in the Facets of Algebraic volume in honor of William Fulton on the occasion of his 80th birthday, <https://arxiv.org/abs/1910.01658>. [↑2](#)
- [Fak12] Najmuddin Fakhruddin, *Chern classes of conformal blocks*, *Compact moduli spaces and vector bundles*, *Contemp. Math.*, vol. 564, Amer. Math. Soc., Providence, RI, 2012, pp. 145–176, DOI 10.1090/conm/564/11148. MR2894632 [↑2](#)
- [Fal94] Gerd Faltings, *A proof for the Verlinde formula*, *J. Algebraic Geom.* **3** (1994), no. 2, 347–374. [↑3](#)
- [FG03] Gavril Farkas and Angela Gibney, *The Mori cones of moduli spaces of pointed curves of small genus*, *Trans. Amer. Math. Soc.* **355** (2003), no. 3, 1183–1199 (electronic). [↑4](#)
- [FBZ04] Edward Frenkel and David Ben-Zvi, *Vertex algebras and algebraic curves*, 2nd ed., *Mathematical Surveys and Monographs*, vol. 88, American Mathematical Society, Providence, RI, 2004. [↑1](#)
- [Gia13] Noah Giansiracusa, *Conformal blocks and rational normal curves*, *J. Algebraic Geom.* **22** (2013), no. 4, 773–793, DOI 10.1090/S1056-3911-2013-00601-3. MR3084722 [↑2](#)
- [GG12] Noah Giansiracusa and Angela Gibney, *The cone of type A, level 1, conformal blocks divisors*, *Adv. Math.* **231** (2012), no. 2, 798–814, DOI 10.1016/j.aim.2012.05.017. MR2955192 [↑2](#), [3](#)
- [GKM02] Angela Gibney, Sean Keel, and Ian Morrison, *Towards the ample cone of  $\overline{M}_{g,n}$* , *J. Amer. Math. Soc.* **15** (2002), no. 2, 273–294. [↑4](#)
- [GM10] Angela Gibney and Diane Maclagan, *Equations for Chow and Hilbert quotients*, *Algebra Number Theory* **4** (2010), no. 7, 855–885. [↑4](#)
- [GJMS13] Angela Gibney, David Jensen, Han-Bom Moon, and David Swinarski, *Veronese quotient models of  $\overline{M}_{0,n}$  and conformal blocks*, *Michigan Math. J.* **62** (2013), no. 4, 721–751. [↑3](#)
- [Gib09] Angela Gibney, *Num criteria for divisors on  $\overline{M}_g$* , *Compos. Math.* **145** (2009), no. 5, 1227–1248. [↑4](#)
- [GM16] Angela Gibney and Swarnava Mukhopadhyay, *On higher Chern classes of vector bundles of conformal blocks*, arXiv:1609.04887 [math.AG] (2016). [↑3](#)
- [GM12] Angela Gibney and Diane Maclagan, *Lower and upper bounds for nef cones*, *Int. Math. Res. Not. IMRN* **14** (2012), 3224–3255. [↑4](#)
- [Gib12] Angela Gibney, *On extensions of the Torelli map*, *Geometry and arithmetic*, EMS Ser. Congr. Rep., Eur. Math. Soc., Zürich, 2012, pp. 125–136. [↑3](#)
- [Gui20] Bin Gui, *Convergence of Sewing Conformal Blocks* (2020). arXiv:2011.07450 [math.QA]. [↑2](#)
- [Hob19] Natalie L. F. Hobson, *Quantum Kostka and the rank one problem for  $\mathfrak{sl}_{2m}$* , *Adv. Geo.* **19** (2019), no. 1, 71–88. [↑3](#)
- [Hua05a] Yi-Zhi Huang, *Diff. equations, duality and modular invariance*, *Comm. Cont. Math.* **7** (2005), no. 5, 649–706. [↑1](#)
- [Hua05b] ———, *Diff. equations and intertwining operators*, *Commun. Contemp. Math.* **7** (2005), no. 3, 375–400. [↑1](#)
- [Hua05c] ———, *Vertex operator algebras, the Verlinde conjecture, and modular tensor categories*, *Proc. Natl. Acad. Sci. USA* **102** (2005), no. 15, 5352–5356. [↑1](#)
- [Hua08] ———, *Rigidity & modularity of vertex tensor categories*, *Commun. Contemp. Math.* **10** (2008), 871–911. [↑1](#)
- [Kap93a] Mikhael Kapranov, *Veronese curves and Grothendieck–Knudsen moduli space  $\overline{M}_{0,n}$* , *J. Algebraic Geom.* **2** (1993), no. 2, 239–262. [↑2](#), [3](#)

- [Kap93b] ———, *Chow quotients of Grassmannians. I*, I. M. Gel'fand Seminar, Adv. Soviet Math., vol. 16, Amer. Math. Soc., Providence, RI, 1993, pp. 29–110. [↑](#)[2](#), [3](#)
- [Kaz16] Anna Kazanova, *On  $S_n$ -invariant conformal blocks vector bundles of rank one on  $\overline{M}_{0,n}$* , Manuscripta Math. **149** (2016), no. 1–2, 107–115. [↑](#)[3](#)
- [KM13] Seán Keel and James McKernan, *Contractible extremal rays on  $\overline{M}_{0,n}$* , Handbook of moduli. Vol. II, Adv. Lect. Math. (ALM), vol. 25, Int. Press, Somerville, MA, 2013, pp. 115–130. [↑](#)[4](#)
- [KNR94] Shrawan Kumar, M. S. Narasimhan, and A. Ramanathan, *Infinite Grassmannians and moduli spaces of  $G$ -bundles*, Math. Ann. **300** (1994), no. 1, 41–75. [↑](#)[3](#)
- [LS97] Yves Laszlo and Christoph Sorger, *The line bundles on the moduli of parabolic  $G$ -bundles over curves and their sections*, Ann. Sci. École Norm. Sup. (4) **30** (1997), no. 4, 499–525. [↑](#)[3](#)
- [MOP15] Alina Marian, Dragos Oprea, and Rahul Pandharipande, *The first Chern class of the Verlinde bundles*, String-Math 2012, Proc. Sympos. Pure Math., vol. 90, Amer. Math. Soc., Providence, RI, 2015, pp. 87–111. [↑](#)[1](#)
- [MOP+17] Alina Marian, Dragos Oprea, Rahul Pandharipande, Aaron Pixton, and Dimitri Zvonkine, *The Chern character of the Verlinde bundle over  $\overline{M}_{g,n}$* , J. Reine Angew. Math. **732** (2017), 147–163. [↑](#)[1](#), [2](#)
- [NT05] Kiyokazu Nagatomo and Akihiro Tsuchiya, *Conformal field theories associated to regular chiral vertex operator algebras. I. Theories over the projective line*, Duke Math. J. **128** (2005), no. 3, 393–471. [↑](#)[1](#)
- [Pau96] Christian Pauly, *Espaces de modules de fibrés parabol et blocs conformes*, Duke **84** (1996), no. 1, 217–235. [↑](#)[3](#)
- [Tsu93] Yoshifumi Tsuchimoto, *Coordinate-free description of conf blocks*, J. Math. Kyoto Univ. **33** (1993), 29–49. [↑](#)[1](#)
- [TK88] Akihiro Tsuchiya and Yukihiro Kanie, *Vertex operators in conformal field theory on  $\mathbf{P}^1$  and monodromy representations of braid group*, Conformal field theory and solvable lattice models (Kyoto, 1986), Adv. Stud. Pure Math., vol. 16, Academic Press, Boston, MA, 1988, pp. 297–372. [↑](#)[1](#)
- [TUY89] Akihiro Tsuchiya, Kenji Ueno, and Yasuhiko Yamada, *Conformal field theory on universal family of stable curves with gauge symmetries*, Integrable systems in quantum field theory and statistical mechanics, Adv. Stud. Pure Math., vol. 19, Academic Press, Boston, MA, 1989, pp. 459–566. [↑](#)[1](#)
- [Zhu94] Yongchang Zhu, *Global vertex ops on Riemann surfaces*, Comm. Math. Phys. **165** (1994), no. 3, 485–531. [↑](#)[1](#)